

Classical Analysis – Math 108b

Homework # 5

due 5 pm Tuesday, February 21, 2006.

1.(10) Exercise 18.57 in [Carothers].

2.(10) Exercise 18.58 in [Carothers].

3.(10) Exercise 18.59 in [Carothers].

4.(10) Let $\{g_n\}$ be a sequence of nonnegative integrable functions possessing the properties

$$i) \lim_{n \rightarrow \infty} g_n(x) = g(x) \text{ exists at a.e. } x \in \mathbb{R}, \quad ii) \lim_{n \rightarrow \infty} \int_{\mathbb{R}} g_n = \int_{\mathbb{R}} g.$$

Suppose another sequence of functions $\{f_n\}$ satisfies $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ almost everywhere and also $|f_n(x)| \leq g_n(x)$ for each n .

Prove that $\lim_{n \rightarrow \infty} \int_{\mathbb{R}} f_n = \int_{\mathbb{R}} f$.

5.(10) Let $p, q, r > 1$ be any choice of real numbers such that $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$. Prove the following extension of Hölder's Inequality:

$$\int_{\mathbb{R}} |f(x)g(x)h(x)| \leq \|f\|_{L_p} \|g\|_{L_q} \|h\|_{L_r}$$

Complementary problems:

6.(5) Let f be a function in $L_p[0, 1]$ for all $p \in [1, P]$. Show that $\|f\|_{L_p}$ is a continuous function of p over the interval $1 \leq p \leq P$. Consider also the case $P = \infty$.

7.(5) Recall that the convolution of two functions f, g is defined as $f * g(x) = \int_{\mathbb{R}} f(x - y)g(y)dy$. Suppose that $f \in L_1(\mathbb{R})$ and $g \in L_{\infty}(\mathbb{R})$. Show that $f * g$ is a bounded and continuous function.