

SOME REMARKS CONCERNING VOEVODSKY'S NILPOTENCE CONJECTURE

MARCELLO BERNARDARA, MATILDE MARCOLLI AND GONÇALO TABUADA

ABSTRACT. In this article we extend Voevodsky's nilpotence conjecture from smooth projective schemes to the broader setting of smooth proper dg categories. Making use of this noncommutative generalization, we then address Voevodsky's original conjecture in the following cases: quadric fibrations, intersection of quadrics, linear sections of Grassmannians, linear sections of determinantal varieties, homological projective duals, and Moizhezon manifolds.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let k be a base field and F a field of coefficients of characteristic zero.

Voevodsky's nilpotence conjecture. In a foundational work [36], Voevodsky introduced the smash-nilpotence equivalence relation $\sim_{\otimes\text{nil}}$ on algebraic cycles and conjectured its agreement with the classical numerical equivalence relation \sim_{num} . Concretely, given a smooth projective k -scheme X , he stated the following:

$$\text{Conjecture } V(X): \mathcal{Z}^*(X)_F / \sim_{\otimes\text{nil}} = \mathcal{Z}^*(X)_F / \sim_{\text{num}}.$$

Thanks to the work of Kahn-Sebastian, Matsusaka, Voevodsky, and Voisin (see [15, 30, 36, 37] and also [1, §11.5.2.3]), the above conjecture holds in the case of curves, surfaces, and abelian 3-folds (when k is of characteristic zero).

Noncommutative nilpotence conjecture. A dg category \mathcal{A} is a category enriched over dg k -vector spaces; see §2.1. Following Kontsevich [17, 18, 19], \mathcal{A} is called *smooth* if it is perfect as a bimodule over itself and *proper* if for any two objects $x, y \in \mathcal{A}$ we have $\sum_i \dim H^i \mathcal{A}(x, y) < \infty$. The classical example is the unique dg enhancement $\text{perf}_{\text{dg}}(X)$ of the category of perfect complexes $\text{perf}(X)$ of a smooth projective k -scheme X ; see Lunts-Orlov [25]. As explained in §2.3-2.4, the Grothendieck group $K_0(\mathcal{A})$ of every smooth proper dg category \mathcal{A} comes endowed with a \otimes -nilpotence equivalence relation $\sim_{\otimes\text{nil}}$ and with a numerical equivalence relation \sim_{num} . Motivated by the above conjecture, we state the following:

$$\text{Conjecture } V_{NC}(\mathcal{A}): K_0(\mathcal{A})_F / \sim_{\otimes\text{nil}} = K_0(\mathcal{A})_F / \sim_{\text{num}}.$$

Our first main result is the following reformulation of Voevodsky's conjecture:

Theorem 1.1. *Conjecture $V(X)$ is equivalent to conjecture $V_{NC}(\text{perf}_{\text{dg}}(X))$.*

Date: November 5, 2014.

2000 Mathematics Subject Classification. 14A22, 14C15, 14D06, 14M10, 14M15, 18G55.

Key words and phrases. Voevodsky's nilpotence conjecture, quadric fibrations, complete intersections, Grassmannians, determinantal varieties, homological projective duality, Moizhezon manifolds, noncommutative algebraic geometry.

Sections 3-5 are a revised version of half of an old preprint [29]. The remaining work is a recent collaboration between the first and the last author.

Theorem 1.1 shows us that when restricted to the commutative world, the noncommutative nilpotence conjecture reduces to Voevodsky’s original conjecture. Making use of this noncommutative viewpoint, we now address Voevodsky’s nilpotence conjecture in several cases.

Quadric fibrations. Let S be a smooth projective k -scheme and $q : Q \rightarrow S$ a flat quadric fibration of relative dimension n with Q smooth. Recall from Kuznetsov [21] (see also [5]) the construction of the sheaf \mathcal{C}_0 of even parts of the Clifford algebra associated to q . Recall also from *loc. cit.* that when the discriminant divisor of q is smooth and n is even (resp. odd) we have a discriminant double cover $\tilde{S} \rightarrow S$ (resp. a square root stack \hat{S}) equipped with an Azumaya algebra \mathcal{B}_0 . Our second main result allows us to decompose conjecture $V(Q)$ into simpler pieces:

Theorem 1.2. *The following holds:*

- (i) *We have $V(Q) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(S, \mathcal{C}_0)) + V(S)$.*
- (ii) *When the discriminant divisor of q is smooth and n is even, we have $V(Q) \Leftrightarrow V(\tilde{S}) + V(S)$. As a consequence, $V(Q)$ holds when $\dim(S) \leq 2$, and becomes equivalent to $V(\tilde{S})$ when S is an abelian 3-fold and k is of characteristic zero.*
- (iii) *When the discriminant divisor of q is smooth and n is odd, we have $V(Q) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(\hat{S}, \mathcal{B}_0)) + V(S)$. As a consequence, $V(Q)$ becomes equivalent to $V_{NC}(\text{perf}_{\text{dg}}(\hat{S}, \mathcal{B}_0))$ when $\dim(S) \leq 2$. This latter conjecture holds when $\dim(S) \leq 2$.*

Remark 1.3. The (rational) Chow motive of a quadric fibration $q : Q \rightarrow S$ was computed by Vial in [35, Thm. 4.2 and Cor. 4.4]. In the particular case where $\dim(S) \leq 2$, it consists of a direct sum of submotives of smooth projective k -schemes of dimension at most two. This motivic decomposition provides an alternative “geometric” proof of conjecture $V(Q)$. We will rely on this argument to prove the last statement of item (iii). In the particular case where S is a curve and k is algebraically closed, we provide also a “categorical” proof of this last statement; see Remark 6.5. The fact that $V_{NC}(\text{perf}_{\text{dg}}(\hat{S}, \mathcal{B}_0))$ holds when $\dim(S) \leq 2$ will play a key role in the proof of Theorem 1.4 below.

Intersection of quadrics. Let X be a smooth complete intersection of r quadric hypersurfaces in \mathbb{P}^m . The linear span of these r quadrics gives rise to a hypersurface $Q \subset \mathbb{P}^{r-1} \times \mathbb{P}^m$, and the projection into the first factor to a flat quadric fibration $q : Q \rightarrow \mathbb{P}^{r-1}$ of relative dimension $m - 1$.

Theorem 1.4. *The following holds:*

- (i) *We have $V(X) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0))$.*
- (ii) *When the discriminant divisor of q is smooth and m is odd, we have $V(X) \Leftrightarrow V(\widehat{\mathbb{P}^{r-1}})$. As a consequence, $V(X)$ holds when $r \leq 3$.*
- (iii) *When the discriminant divisor of q is smooth and m is even, we have $V(X) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(\widehat{\mathbb{P}^{r-1}}, \mathcal{B}_0))$. This latter conjecture holds when $r \leq 3$ and k is algebraically closed.*

Remark 1.5. The (rational) Chow motive of a complete intersection X was computed in [8, Thm. 2.1] in the particular cases where $r \leq 2$ or $r = 3$ and m is odd. It consists of a direct sum of submotives of smooth projective k -schemes of dimension

at most one. This motivic decomposition provides an alternative proof of conjecture $V(X)$. A similar argument holds in the case where $r = 3$ and m is even.

Remark 1.6. (Relative version) Theorem 1.4 has a relative analogue with X replaced by a generic relative complete intersection $X \rightarrow S$ of r quadric fibrations $Q_i \rightarrow S$ of relative dimension $m - 1$; consult [5, Def. 1.2.4] for details. Items (i), (ii), and (iii), hold similarly with \mathbb{P}^{r-1} replaced by a \mathbb{P}^{r-1} -bundle $T \rightarrow S$, with $V(\widehat{\mathbb{P}^{r-1}})$ replaced by $V(\widehat{T}) + V(S)$, and with $V_{NC}(\text{perf}_{\text{dg}}(\widehat{\mathbb{P}^{r-1}}, \mathcal{B}_0))$ replaced by $V_{NC}(\text{perf}_{\text{dg}}(\widehat{T}, \mathcal{B}_0)) + V(S)$, respectively. Note that thanks to the relative item (ii), conjecture $V(X)$ holds when $r = 2$ and S is a curve.

Linear sections of Grassmanians. Following Kuznetsov [22], consider the following two classes of schemes:

- (i) Let X_L be a generic linear section of codimension r of the Grassmannian $\text{Gr}(2, W)$ (with $W = k^{\oplus 6}$) under the Plücker embedding, and Y_L the corresponding dual linear section of the cubic Pfaffian $\text{Pf}(4, W^*)$ in $\mathbb{P}(\Lambda^2 W^*)$.

For example when $r = 3$, X_L is a Fano 5-fold; when $r = 4$, X_L is a Fano 4-fold; and when $r = 6$, X_L is a $K3$ surface of degree 14 and Y_L a Pfaffian cubic 4-fold. Moreover, X_L and Y_L are smooth whenever $r \leq 6$.

- (ii) Let X_L be a generic linear section of codimension r of the Grassmannian $\text{Gr}(2, W)$ (with $W = k^{\oplus 7}$) under the Plücker embedding, and Y_L the corresponding dual linear section of the cubic Pfaffian $\text{Pf}(4, W^*)$ in $\mathbb{P}(\Lambda^2 W^*)$.

For example when $r = 5$, X_L is a Fano 5-fold; when $r = 4$, X_L is a Fano 4-fold; when $r = 8$, Y_L is a Fano 4-fold; and when $r = 9$, Y_L is a Fano 5-fold. Moreover, X_L and Y_L are smooth whenever $r \leq 10$.

Theorem 1.7. *Let X_L and Y_L be as in the above classes (i)-(ii). Under the assumption that X_L and Y_L are smooth, we have $V(X_L) \Leftrightarrow V(Y_L)$. This conjecture holds when $r \leq 6$ (class (i)), and when $r \leq 6$ and $8 \leq r \leq 10$ (class (ii)).*

Remark 1.8. To the best of the authors' knowledge, Theorem 1.7 proves Voevodsky's nilpotence conjecture in new cases.

Linear sections of determinantal varieties. Let U and V be k -vector spaces of dimensions m and n , respectively, with $n \geq m$, and $0 < r < m$ an integer. Following [7], consider the universal determinantal variety $Z_{m,n}^r \subset \mathbb{P}(U \otimes V)$ given by the locus of matrices $M : U \rightarrow V^*$ of rank at most r . Its Springer resolution is denoted by $\mathcal{X}_{m,n}^r := \mathbb{P}(\mathcal{Q} \otimes U) \rightarrow \text{Gr}(r, U)$, where \mathcal{Q} stands for the tautological quotient on $\text{Gr}(r, U)$. Under these notations, we have the following class of schemes:

- (i) Let X_L be a generic linear section of codimension c of $\mathcal{X}_{m,n}^r$ under the map $\mathcal{X}_{m,n}^r \rightarrow \mathbb{P}(U \otimes V)$, and Y_L the corresponding dual linear section of $\mathcal{X}_{m,n}^{m-r}$ under the map $\mathcal{X}_{m,n}^{m-r} \rightarrow \mathbb{P}(U^* \otimes V^*)$.

Remark 1.9. As explained in [7, §3], X_L and Y_L are smooth crepant categorical resolution of singularities of $(Z_{m,n}^r)_L$ and $(Z_{m,n}^{m-r})_L$, respectively.

For example when $m = n = 4$ and $r = 1$, X_L is a $(6 - c)$ -dimensional section of $\mathbb{P}^3 \times \mathbb{P}^3$ under the Segre embedding, and Y_L the resolution of the dual $(c - 2)$ -dimensional determinantal quartic. In the same vein, when $m = n = 4$ and $r = 2$, X_L is a $(11 - c)$ -dimensional section of the self-dual orbit of 4×4 matrices of rank 2, and Y_L the $(c - 5)$ -dimensional dual section. Moreover, the following holds:

- (a) When $c \leq 7$, X_L is a Fano $(11 - 4)$ -fold and $\dim(Y_L) \leq 2$.
- (b) When $c = 8$, X_L and Y_L are dual Calabi-Yau threefolds
- (c) When $c \geq 9$, Y_L is a Fano $(c - 5)$ -fold and $\dim(X_L) \leq 2$.

For further example, consult [7, §3.3] and well as Tables 1 and 2 in *loc. cit.*

Theorem 1.10. *When X_L and Y_L are as in the above class (i), we have $V(X_L) \Leftrightarrow V(Y_L)$. This conjecture holds when $\dim(X_L) \leq 2$ or $\dim(Y_L) \leq 2$.*

Remark 1.11. To the best of the authors' knowledge, Theorem 1.10 proves Voevodsky's nilpotence conjecture in new cases.

Homological projective duality. Making use of Kuznetsov's theory of homological projective duality (HPD) [23], Theorem 1.7 admits the following generalization: let X be a smooth projective k -scheme equipped with an ample line bundle $\mathcal{O}_X(1)$. Note that $\mathcal{O}_X(1)$ gives rise to a morphism $X \rightarrow \mathbb{P}(V)$, where $V := H^0(X, \mathcal{O}_X(1))^*$. Let Y be the HP-dual of X , $\mathcal{O}_Y(1)$ the associated ample line bundle, and $Y \rightarrow \mathbb{P}(V^*)$ the associated morphism. Assume that $\text{perf}(X)$ admits a *Lefschetz decomposition* $\text{perf}(X) = \langle \mathbb{A}_0, \dots, \mathbb{A}_n(n) \rangle$ with respect to $\mathcal{O}_X(1)$, i.e. a semi-orthogonal decomposition of $\text{perf}(X)$ such that $\mathbb{A}_0 \supset \dots \supset \mathbb{A}_n$ and $\mathbb{A}_i(i) := \mathbb{A}_i \otimes \mathcal{O}(i)$. Assume also that conjecture $V_{NC}(\mathbb{A}_0^{\text{dg}})$ holds, where \mathbb{A}_0^{dg} stands for the dg enhancement of \mathbb{A}_0 induced from $\text{perf}_{\text{dg}}(X)$; see §2.2. Finally, let $L \subset V$ be a subspace such that the linear sections $X_L := X \times_{\mathbb{P}(V)} \mathbb{P}(L^\perp)$ and $Y_L := Y \times_{\mathbb{P}(V^*)} \mathbb{P}(L)$ are of expected dimension $\dim(X_L) = \dim(X) - \dim(L)$ and $\dim(Y_L) = \dim(Y) - \dim(L^\perp)$, respectively.

Theorem 1.12. *Let X_L and Y_L be as above. Under the assumption that X_L and Y_L are smooth, we have $V(X_L) \Leftrightarrow V(Y_L)$.*

Remark 1.13. Theorem 1.12 reduces to Theorem 1.7 (resp. to Theorem 1.10) in the particular case of Grassmanian-Pfaffian (resp. determinantal) homological projective duality; consult [22] (resp. [7]) for details.

Moishezon manifolds. A *Moishezon manifold* X is a compact complex manifold such that the field of meromorphic functions on each component of X has transcendence degree equal to the dimension of the component. As proved by Moishezon [31], X is a smooth projective \mathbb{C} -scheme if and only if it admits a Kähler metric. In the remaining cases, Artin [3] showed that X is a proper algebraic space over \mathbb{C} .

Let $Y \rightarrow \mathbb{P}^2$ be one of the non-rational conic bundles described by Artin and Mumford in [4], and $X \rightarrow Y$ a small resolution. In this case, X is a smooth (non necessarily projective) Moishezon manifold.

Theorem 1.14. *Conjecture $V_{NC}(\text{perf}_{\text{dg}}(X))$ holds for the above resolutions.*

Remark 1.15. The proofs of Theorems 1.2, 1.4, 1.7, 1.10, 1.12, and 1.14 are based on the study of a smooth projective k -scheme (or algebraic space) X via semi-orthogonal decompositions of its category of perfect complexes $\text{perf}(X)$; see Bondal-Orlov [9] and Kuznetsov [24] for instance. This approach allows the reduction of Voevodsky's conjecture $V(X)$ to several noncommutative conjectures V_{NC} - one for each piece of the semi-orthogonal decomposition. We believe this provides a new tool for the proof of Voevodsky's conjecture as well as of its generalization to algebraic spaces.

2. PRELIMINARIES

2.1. Dg categories. A *differential graded (=dg) category* \mathcal{A} is a category enriched over dg k -vector spaces; consult Keller [16] for details. For example, every (dg) k -algebra A gives naturally rise to a dg category \underline{A} with a single object. Let dgcats be the category of small dg categories. Recall from [16, §3] the construction of the derived category $\mathcal{D}(\mathcal{A})$ of \mathcal{A} . This triangulated category admits arbitrary direct sums and we will write $\mathcal{D}_c(\mathcal{A})$ for the full subcategory of compact objects. A dg functor $\mathcal{A} \rightarrow \mathcal{B}$ is called a *Morita equivalence* if it induces an equivalence $\mathcal{D}(\mathcal{A}) \xrightarrow{\sim} \mathcal{D}(\mathcal{B})$. Finally, let us write $\mathcal{A} \otimes \mathcal{B}$ for the tensor product of dg categories.

2.2. Perfect complexes. Given a stack \mathcal{X} and a sheaf of $\mathcal{O}_{\mathcal{X}}$ -algebras \mathcal{G} , let $\text{Mod}(\mathcal{X}, \mathcal{G})$ be the Grothendieck category of sheaves of (right) \mathcal{G} -modules, $\mathcal{D}(\mathcal{X}, \mathcal{G}) := \mathcal{D}(\text{Mod}(\mathcal{X}, \mathcal{G}))$ the derived category of \mathcal{G} , and $\text{perf}(\mathcal{X}, \mathcal{G})$ the subcategory of perfect complexes. As explained in [16, §4.4], the derived category $\mathcal{D}_{\text{dg}}(\mathcal{E}x)$ of an abelian (or exact) category $\mathcal{E}x$ is defined as the (Drinfeld's) dg quotient $\mathcal{C}_{\text{dg}}(\mathcal{E}x)/\text{Ac}_{\text{dg}}(\mathcal{E}x)$ of the dg category of complexes over $\mathcal{E}x$ by its full dg subcategory of acyclic complexes. Hence, let us write $\mathcal{D}_{\text{dg}}(\mathcal{X}, \mathcal{G})$ for the dg category $\mathcal{D}_{\text{dg}}(\mathcal{E}x)$ with $\mathcal{E}x := \text{Mod}(\mathcal{X}, \mathcal{G})$ and $\text{perf}_{\text{dg}}(\mathcal{X}, \mathcal{G})$ for the full dg subcategory of perfect complexes.

Lemma 2.1. *Let X be a smooth projective k -scheme and $\text{perf}(X) = \langle \mathcal{T}_1, \dots, \mathcal{T}_n \rangle$ a semi-orthogonal decomposition. In this case, the dg categories $\mathcal{T}_i^{\text{dg}}$ (where $\mathcal{T}_i^{\text{dg}}$ stands for the dg enhancement of \mathcal{T}_i induced from $\text{perf}_{\text{dg}}(X)$) are smooth and proper.*

Proof. Let $\text{Ho}(\text{dgcats})$ be the localization of dgcats with respect to the class of Morita equivalences. The tensor product of dg categories gives rise to a symmetric monoidal structure on dgcats which descends to $\text{Ho}(\text{dgcats})$. Moreover, as proved in [10, Thm. 5.8], the smooth and proper dg categories can be characterized as those objects of $\text{Ho}(\text{dgcats})$ which are dualizable. Note that the canonical inclusion $\mathcal{T}_i^{\text{dg}} \hookrightarrow \text{perf}_{\text{dg}}(X)$ and projection $\text{perf}_{\text{dg}}(X) \rightarrow \mathcal{T}_i^{\text{dg}}$ dg functors express $\mathcal{T}_i^{\text{dg}}$ as a direct factor of $\text{perf}_{\text{dg}}(X)$ in $\text{Ho}(\text{dgcats})$. Hence, since $\text{perf}_{\text{dg}}(X)$ is smooth and proper, we conclude that $\mathcal{T}_i^{\text{dg}}$ is also smooth and proper. \square

2.3. \otimes -nilpotence equivalence relation. Let \mathcal{A} be a dg category. An element $[M]$ of the Grothendieck group $K_0(\mathcal{A}) := K_0(\mathcal{D}_c(\mathcal{A}))$ is called *\otimes -nilpotent* if there exists an integer $n > 0$ such that $[M^{\otimes n}] = 0$ in $K_0(\mathcal{A}^{\otimes n})$. This gives rise to a well-defined equivalence relation $\sim_{\otimes\text{nil}}$ on $K_0(\mathcal{A})$ and on its F -linearization $K_0(\mathcal{A})_F$.

2.4. Numerical equivalence relation. Let \mathcal{A} be a smooth proper dg category. As explained in [27, §4], the pairing $(M, N) \mapsto \sum_i (-1)^i \dim \text{Hom}_{\mathcal{D}_c(\mathcal{A})}(M, N[i])$ gives rise to a well-defined bilinear form $\chi(-, -)$ on $K_0(\mathcal{A})$. Moreover, the left and right kernels of $\chi(-, -)$ are the same. An element $[M]$ of the Grothendieck group $K_0(\mathcal{A})$ is called *numerically trivial* if $\chi([M], [N]) = 0$ for all $[N] \in K_0(\mathcal{A})$. This gives rise to an equivalence relation \sim_{num} on $K_0(\mathcal{A})$ and consequently on $K_0(\mathcal{A})_F$. When $\mathcal{A} = \text{perf}_{\text{dg}}(X)$, with X a smooth projective k -scheme, and $F = \mathbb{Q}$ this equivalence relation reduces, via the Chern character $K_0(X)_{\mathbb{Q}} \xrightarrow{\sim} CH^*(X)_{\mathbb{Q}}$, to the classical numerical equivalence relation on the Chow ring $CH^*(X)_{\mathbb{Q}}$.

2.5. Motives. We assume the reader is familiar with the categories of Chow motives $\text{Chow}(k)_F$ and numerical motives $\text{Num}(k)_F$; see [1, §4]. The Tate motive will be denoted $F(1)$. In the same vein, we assume some familiarity with the categories

of noncommutative Chow motives $\mathrm{NChow}(k)_F$ and noncommutative numerical motives $\mathrm{NNum}(k)_F$; consult the surveys [28, 2-3] [33, 4] and the references therein. Recall from *loc. cit.* that $\mathrm{NNum}(k)_F$ is the idempotent completion of the quotient of $\mathrm{NChow}(k)_F$ by its largest \otimes -ideal¹, and that $\mathrm{Hom}_{\mathrm{NChow}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F$.

3. ORBIT CATEGORIES AND \otimes -NILPOTENCE

Let \mathcal{C} be an F -linear additive rigid symmetric monoidal category.

Orbit categories. Given a \otimes -invertible object $\mathcal{O} \in \mathcal{C}$, recall from [32, 7] the construction of the orbit category $\mathcal{C}/_{-\otimes\mathcal{O}}$. It has the same objects as \mathcal{C} and morphisms

$$\mathrm{Hom}_{\mathcal{C}/_{-\otimes\mathcal{O}}}(a, b) := \bigoplus_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(a, b \otimes \mathcal{O}^{\otimes j}).$$

The composition law is induced from \mathcal{C} . By construction, $\mathcal{C}/_{-\otimes\mathcal{O}}$ is F -linear, additive, and comes equipped with a canonical projection functor $\pi : \mathcal{C} \rightarrow \mathcal{C}/_{-\otimes\mathcal{O}}$. Moreover, π is endowed with a natural 2-isomorphism $\pi \circ (- \otimes \mathcal{O}) \xrightarrow{\cong} \pi$ and is 2-universal among all such functors. As proved in [32, Lem. 7.3], $\mathcal{C}/_{-\otimes\mathcal{O}}$ inherits from \mathcal{C} a symmetric monoidal structure making π symmetric monoidal. On objects it is the same. On morphisms it is defined as the unique bilinear pairing

$$\bigoplus_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(a, b \otimes \mathcal{O}^{\otimes j}) \times \bigoplus_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(c, d \otimes \mathcal{O}^{\otimes j}) \longrightarrow \bigoplus_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(a \otimes c, (b \otimes d) \otimes \mathcal{O}^{\otimes j})$$

which sends the pair $(a \xrightarrow{f_r} b \otimes \mathcal{O}^{\otimes r}, c \xrightarrow{g_s} d \otimes \mathcal{O}^{\otimes s})$ to

$$(f \otimes g)_{(r+s)} : a \otimes c \xrightarrow{f_r \otimes g_s} b \otimes \mathcal{O}^{\otimes r} \otimes d \otimes \mathcal{O}^{\otimes s} \simeq (b \otimes d) \otimes \mathcal{O}^{\otimes(r+s)}.$$

\otimes -nilpotence. The \otimes_{nil} -ideal of \mathcal{C} is defined as

$$\otimes_{\mathrm{nil}}(a, b) := \{f \in \mathrm{Hom}_{\mathcal{C}}(a, b) \mid f^{\otimes n} = 0 \text{ for } n \gg 0\}.$$

By construction, \otimes_{nil} is a \otimes -ideal. Moreover, all its ideals $\otimes_{\mathrm{nil}}(a, a) \subset \mathrm{Hom}_{\mathcal{C}}(a, a)$ are nilpotent; see [2, Lem. 7.4.2 (ii)]. As a consequence, the \otimes -functor $\mathcal{C} \rightarrow \mathcal{C}/_{\otimes_{\mathrm{nil}}}$ is not only F -linear and additive but moreover conservative. Furthermore, since idempotents can be lifted along nilpotent ideals (see [6, III Prop. 2.10]), $\mathcal{C}/_{\otimes_{\mathrm{nil}}}$ is idempotent complete whenever \mathcal{C} is idempotent complete.

Compatibility. Let \mathcal{C} be a category and $\mathcal{O} \in \mathcal{C}$ a \otimes -invertible objects as above.

Proposition 3.1. *There exists a canonical F -linear additive \otimes -equivalence θ making the following diagram commute:*

$$(3.2) \quad \begin{array}{ccc} \mathcal{C}/_{\otimes_{\mathrm{nil}}} & \longleftarrow \mathcal{C} & \longrightarrow \mathcal{C}/_{-\otimes\mathcal{O}} \\ \downarrow & & \downarrow \\ (\mathcal{C}/_{\otimes_{\mathrm{nil}}})/_{-\otimes\mathcal{O}} & \xrightarrow[\theta]{\cong} & (\mathcal{C}/_{-\otimes\mathcal{O}})/_{\otimes_{\mathrm{nil}}} \end{array}.$$

Proof. The existence of the F -linear additive \otimes -functor θ follows from the fact that

$$(3.3) \quad \mathcal{C} \longrightarrow \mathcal{C}/_{-\otimes\mathcal{O}} \longrightarrow (\mathcal{C}/_{-\otimes\mathcal{O}})/_{\otimes_{\mathrm{nil}}}$$

vanishes on the \otimes_{nil} -ideal and also from the natural 2-isomorphism between (3.3) $\circ (- \otimes \mathcal{O})$ and (3.3). Note that the functor θ is the identity on objects and sends $\{[f_j]\}_{j \in \mathbb{Z}}$ to $\{[f_j]\}_{j \in \mathbb{Z}}$. Clearly, it is full. The faithfulness is left as an exercise. \square

¹Different from the entire category $\mathrm{NChow}(k)_F$.

4. \otimes -NILPOTENCE OF MOTIVES

By construction, the categories $\text{Chow}(k)_F$ and $\text{NChow}(k)_F$ are F -linear, additive, rigid symmetric monoidal, and idempotent complete. Let us denote by

$$\text{Voev}(k)_F := \text{Chow}(k)_F / \otimes_{\text{nil}} \quad \text{and} \quad \text{NVoev}(k)_F := \text{NChow}(k)_F / \otimes_{\text{nil}}$$

the associated quotients. They fit in the following sequences

$$\text{Chow}(k)_F \rightarrow \text{Voev}(k)_F \rightarrow \text{Num}(k)_F \quad \text{NChow}(k)_F \rightarrow \text{NVoev}(k)_F \rightarrow \text{NNum}(k)_F .$$

The relation between all these motivic categories is the following:

Proposition 4.1. *There exist F -linear additive fully-faithful \otimes -functors $R, R_{\otimes_{\text{nil}}}, R_{\mathcal{N}}$ making the following diagram commute:*

$$(4.2) \quad \begin{array}{ccccc} \text{Chow}(k)_F & \xrightarrow{\pi} & \text{Chow}(k)_{F/-\otimes F(1)} & \xrightarrow{R} & \text{NChow}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Voev}(k)_F & \xrightarrow{\pi} & \text{Voev}(k)_{F/-\otimes F(1)} & \xrightarrow{R_{\otimes_{\text{nil}}}} & \text{NVoev}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Num}(k)_F & \xrightarrow{\pi} & \text{Num}(k)_{F/-\otimes F(1)} & \xrightarrow{R_{\mathcal{N}}} & \text{NNum}(k)_F . \end{array}$$

Proof. The outer commutative square, with $R, R_{\mathcal{N}}$ F -linear additive fully-faithful \otimes -functors, was built in [26, Thm. 1.13]. Consider now the ‘‘zoomed’’ diagram:

$$(4.3) \quad \begin{array}{ccccc} \text{Chow}(k)_{F/-\otimes F(1)} & \xlongequal{\quad} & \text{Chow}(k)_{F/-\otimes F(1)} & \xrightarrow{R} & \text{NChow}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Voev}(k)_{F/-\otimes F(1)} & \xrightarrow[\theta]{\simeq} & (\text{Chow}(k)_{F/-\otimes F(1)}) / \otimes_{\text{nil}} & \xrightarrow{R / \otimes_{\text{nil}}} & \text{NVoev}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Num}(k)_{F/-\otimes F(1)} & \xlongequal{\quad} & \text{Num}(k)_{F/-\otimes F(1)} & \xrightarrow{R_{\mathcal{N}}} & \text{NNum}(k)_F . \end{array}$$

By definition, $R_{\otimes_{\text{nil}}} := R / \otimes_{\text{nil}} \circ \theta$. Hence, since R is an F -linear additive fully-faithful \otimes -functor, we conclude that $R_{\otimes_{\text{nil}}}$ is also an F -linear additive fully-faithful \otimes -functor. The commutativity of the bottom squares of diagram (4.3) follows from the fact that $\text{Num}(k)_{F/-\otimes F(1)}$ identifies with the quotient of $\text{Chow}(k)_{F/-\otimes F(1)}$ by its largest \otimes -ideal \mathcal{N} ; consult [26, Prop. 3.2] for details. \square

5. PROOF OF THEOREM 1.1

Note first that we have the following natural isomorphisms

$$\begin{aligned} \text{Hom}_{\text{Voev}(k)_{F/-\otimes F(1)}}(\text{Spec}(k), X) &\simeq \mathcal{Z}^*(X)_F / \sim_{\otimes_{\text{nil}}} \\ \text{Hom}_{\text{Num}(k)_{F/-\otimes F(1)}}(\text{Spec}(k), X) &\simeq \mathcal{Z}^*(X)_F / \sim_{\text{num}} . \end{aligned}$$

As a consequence, conjecture $V(X)$ becomes equivalent to the injectivity of

$$(5.1) \quad \text{Hom}_{\text{Voev}(k)_{F/-\otimes F(1)}}(\text{Spec}(k), X) \twoheadrightarrow \text{Hom}_{\text{Num}(k)_{F/-\otimes F(1)}}(\text{Spec}(k), X) .$$

Given a smooth and proper dg category \mathcal{A} , we have also natural isomorphisms

$$\text{Hom}_{\text{NVoev}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F / \sim_{\otimes_{\text{nil}}} \quad \text{Hom}_{\text{NNum}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F / \sim_{\text{num}} .$$

Hence, conjecture $V_{NC}(\mathcal{A})$ becomes equivalent to the injectivity of

$$(5.2) \quad \mathrm{Hom}_{\mathrm{NVoev}(k)_F}(\underline{k}, \mathcal{A}) \rightarrow \mathrm{Hom}_{\mathrm{NNum}(k)_F}(\underline{k}, \mathcal{A}).$$

Now, recall from [32, Thm. 1.1] that the image of X under the composed functor $R \circ \pi$ identifies naturally with the noncommutative Chow motive $\mathrm{perf}_{\mathrm{dg}}(X)$. Similarly, the image of $\mathrm{Spec}(k)$ under $R \circ \pi$ identifies with $\mathrm{perf}_{\mathrm{dg}}(\mathrm{Spec}(k))$ which is Morita equivalent to \underline{k} . As a consequence, since the functors $R_{\otimes_{\mathrm{nil}}}$ and $R_{\mathcal{N}}$ are fully-faithful, the bottom right-hand side square of diagram (4.2) gives rise to the following commutative diagram:

$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{Voev}(k)_{F/-\otimes F(1)}}(\mathrm{Spec}(k), X) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathrm{NVoev}(k)_F}(\underline{k}, \mathrm{perf}_{\mathrm{dg}}(X)) \\ (5.1) \downarrow & & \downarrow (5.2) \\ \mathrm{Hom}_{\mathrm{Num}(k)_{F/-\otimes F(1)}}(\mathrm{Spec}(k), X) & \xrightarrow{\simeq} & \mathrm{Hom}_{\mathrm{NNum}(k)_F}(\underline{k}, \mathrm{perf}_{\mathrm{dg}}(X)). \end{array}$$

Using the above reformulations of conjectures V and V_{NC} , we conclude finally that conjecture $V(X)$ is equivalent to conjecture $V_{NC}(\mathrm{perf}_{\mathrm{dg}}(X))$.

6. PROOF OF THEOREM 1.2

Item (i). As proved by Kuznetsov in [21, Thm. 4.2], one has the following semi-orthogonal decomposition

$$\mathrm{perf}(Q) = \langle \mathrm{perf}(S, \mathcal{C}_0), \mathrm{perf}(S)_1, \dots, \mathrm{perf}(S)_n \rangle$$

with $\mathrm{perf}(S)_i := q^* \mathrm{perf}(S) \otimes \mathcal{O}_{Q/S}(i)$. Note that $\mathrm{perf}(S)_i \simeq \mathrm{perf}(S)$ for every i . Using [8, Prop. 3.1], one then obtains a direct sum decomposition in $\mathrm{NChow}(k)_F$

$$(6.1) \quad \mathrm{perf}_{\mathrm{dg}}(Q) \simeq \mathrm{perf}^{\mathrm{dg}}(S, \mathcal{C}_0) \oplus \underbrace{\mathrm{perf}_{\mathrm{dg}}(S) \oplus \dots \oplus \mathrm{perf}_{\mathrm{dg}}(S)}_{n \text{ copies}},$$

where $\mathrm{perf}^{\mathrm{dg}}(S, \mathcal{C}_0)$ stands for the dg enhancement of $\mathrm{perf}(S, \mathcal{C}_0)$ induced from $\mathrm{perf}_{\mathrm{dg}}(Q)$. Note that thanks to Lemma 2.1, the dg category $\mathrm{perf}^{\mathrm{dg}}(S, \mathcal{C}_0)$ is smooth and proper. Since the inclusion of categories $\mathrm{perf}(S, \mathcal{C}_0) \hookrightarrow \mathrm{perf}(Q)$ is of Fourier-Mukai type (see [21, Prop. 4.9]), its kernel $\mathcal{K} \in \mathrm{perf}(S \times Q, \mathcal{C}_0^{\mathrm{op}} \boxtimes \mathcal{O}_X)$ gives rise to a Fourier-Mukai Morita equivalence $\Phi_{\mathrm{dg}}^{\mathcal{K}} : \mathrm{perf}_{\mathrm{dg}}(S, \mathcal{C}_0) \rightarrow \mathrm{perf}^{\mathrm{dg}}(S, \mathcal{C}_0)$. Hence, we can replace in the above decomposition (6.1) the dg category $\mathrm{perf}^{\mathrm{dg}}(S, \mathcal{C}_0)$ by the canonical one $\mathrm{perf}_{\mathrm{dg}}(S, \mathcal{C}_0)$ (see §2.2). Finally, using the above description (5.2) of the noncommutative nilpotence conjecture, one concludes that

$$(6.2) \quad V_{NC}(\mathrm{perf}_{\mathrm{dg}}(Q)) \Leftrightarrow V_{NC}(\mathrm{perf}_{\mathrm{dg}}(S, \mathcal{C}_0)) + V_{NC}(\mathrm{perf}_{\mathrm{dg}}(S)).$$

The proof follows now automatically from Theorem 1.1.

Item (ii). As proved by Kuznetsov in [21, Prop. 3.13], $\mathrm{perf}(S, \mathcal{C}_0)$ is Fourier-Mukai equivalent to $\mathrm{perf}(\tilde{S}, \mathcal{B}_0)$. Hence, the above equivalence (6.2) reduces to

$$(6.3) \quad V_{NC}(\mathrm{perf}_{\mathrm{dg}}(Q)) \Leftrightarrow V_{NC}(\mathrm{perf}_{\mathrm{dg}}(\tilde{S}, \mathcal{B}_0)) + V_{NC}(\mathrm{perf}_{\mathrm{dg}}(S)).$$

Since \mathcal{B}_0 is a sheaf of Azumaya algebras and F is of characteristic zero, the canonical dg functor $\mathrm{perf}_{\mathrm{dg}}(\tilde{S}) \rightarrow \mathrm{perf}_{\mathrm{dg}}(\tilde{S}, \mathcal{B}_0)$ becomes an isomorphism in $\mathrm{NChow}(k)_F$; see [34, Thm. 2.1]. Consequently, conjecture $V_{NC}(\mathrm{perf}_{\mathrm{dg}}(\tilde{S}, \mathcal{B}_0))$ reduces to conjecture $V_{NC}(\mathrm{perf}_{\mathrm{dg}}(\tilde{S}))$. The proof follows now from Theorem 1.1.

Item (iii). As proved by Kuznetsov in [21, Prop. 3.15], $\text{perf}(S, \mathcal{C}_0)$ is Fourier-Mukai equivalent to $\text{perf}(\widehat{S}, \mathcal{B}_0)$. Hence, the above equivalence (6.2) reduces to

$$(6.4) \quad V_{NC}(\text{perf}_{\text{dg}}(Q)) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(\widehat{S}, \mathcal{B}_0)) + V_{NC}(\text{perf}_{\text{dg}}(S)).$$

The proof of the first claim follows now from Theorem 1.1.

Let us now prove the second claim, which via (6.4) is equivalent to the proof of $V_{NC}(\text{perf}_{\text{dg}}(Q))$. Thanks to Vial [35, Thm. 4.2 and Cor. 4.4], the rational Chow motive $M_{\mathbb{Q}}(Q)$ of Q decomposes as $M_{\mathbb{Q}}(Q) = M_{\mathbb{Q}}(S)^{\oplus(n-\dim(S))} \oplus N$, where N stands for a submotive of a smooth projective k -scheme of dimension $\leq \dim(S)$. Therefore, when $\dim(S) \leq 2$, conjecture $V(Q) = V_{NC}(\text{perf}_{\text{dg}}(Q))$ holds.

Remark 6.5. Assume that S is a smooth projective curve and that k is algebraically closed. In this remark we provide a ‘‘categorical’’ proof of the second claim of item (iii) of Theorem 1.2. Thanks to the work of Graber-Harris-Starr [13], the fibration $q : Q \rightarrow S$ admits a section. Making use of it, we can perform reduction by hyperbolic splitting in order to obtain a conic bundle $q' : Q' \rightarrow S$; consult [5, §1.3] for details. The sheaf \mathcal{C}'_0 of even parts of the associated Clifford algebra is such that the categories $\text{perf}(S, \mathcal{C}_0)$ and $\text{perf}(S, \mathcal{C}'_0)$ are Fourier-Mukai equivalent; see [5, Rk. 1.8.9]. As a consequence, using (6.2) and the fact that $\dim(S) = 1$, we obtain the following equivalence:

$$(6.6) \quad V_{NC}(\text{perf}_{\text{dg}}(Q)) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(Q')).$$

Since S is a curve, Q' is a surface. Therefore, conjecture (6.6) holds.

7. PROOF OF THEOREM 1.4

Item (i). As proved by Kuznetsov in [21, Thm. 5.5], we have a Fourier-Mukai equivalence $\text{perf}(X) \simeq \text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0)$ when $m - 2r + 1 = 0$, the following semi-orthogonal decomposition

$$\text{perf}(X) = \langle \text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0), \mathcal{O}_X(1), \dots, \mathcal{O}_X(m - 2r + 1) \rangle,$$

when $m - 2r + 1 > 0$, and a dual semi-orthogonal decomposition of $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0)$ (containing a copy of $\text{perf}(X)$ and exceptional objects) when $m - 2r + 1 < 0$. The proof of the case $m - 2r + 1 = 0$ is clear. Let us now prove the case $m - 2r + 1 > 0$; the proof of the case $m - 2r + 1 < 0$ is similar. Using [8, Prop. 3.11], one obtains the following direct sum decomposition in $\text{NChow}(k)_F$

$$\text{perf}_{\text{dg}}(X) \simeq \text{perf}^{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0) \oplus \underbrace{\text{perf}_{\text{dg}}(\underline{k}) \oplus \dots \oplus \text{perf}_{\text{dg}}(\underline{k})}_{(m-2r+1) \text{ copies}}.$$

Thanks to Lemma 2.1, the dg category $\text{perf}^{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0)$ is smooth and proper². Since $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0) \hookrightarrow \text{perf}(X)$ is of Fourier-Mukai type (see [21, Prop. 4.9]), an argument similar to the one of the proof of Theorem 1.2(i) shows us that

$$V_{NC}(\text{perf}_{\text{dg}}(X)) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0)) + V_{NC}(\text{perf}_{\text{dg}}(\underline{k})).$$

The proof follows now automatically from Theorem 1.1.

Item (ii)-(iii). The proofs are similar to those of items (ii)-(iii) of Theorem 1.2.

²In the case $m - 2r + 1 < 0$, these properties follow from the existence of a fully faithful Fourier-Mukai functor $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0) \rightarrow \text{perf}(Q)$, with $Q \subset \mathbb{P}^{r-1} \times \mathbb{P}^m$ a smooth hypersurface.

8. PROOF OF THEOREMS 1.7 AND 1.10

Assume that X_L and Y_L are as in classes of (i)-(ii) of Theorem 1.7 (resp. as in class (i) of Theorem 1.10). As proved by Kuznetsov in [22, §10-11] (resp. in [7, Thm. 3.4]), one of the following three situations occurs:

- (a) there is a semi-orthogonal decomposition $\text{perf}(X_L) = \langle \text{perf}(Y_L), \mathcal{E}_1, \dots, \mathcal{E}_n \rangle$, with the \mathcal{E}_i 's exceptional bundles on X_L ;
- (b) there is a semi-orthogonal decomposition $\text{perf}(Y_L) = \langle \text{perf}(X_L), \mathcal{E}'_1, \dots, \mathcal{E}'_n \rangle$, with the \mathcal{E}'_i 's exceptional bundles on Y_L ;
- (c) there is a Fourier-Mukai equivalence between $\text{perf}(X_L)$ and $\text{perf}(Y_L)$.

Therefore, equivalence $V(X_L) \Leftrightarrow V(Y_L)$ is clear in situation (c). Since the inclusions of categories $\text{perf}(Y_L) \hookrightarrow \text{perf}(X_L)$ (situation (a)) and $\text{perf}(X_L) \hookrightarrow \text{perf}(Y_L)$ (situation (b)) are of Fourier-Mukai type, a proof similar to the one of Theorem 1.2(i) shows us that equivalence $V(X_L) \Leftrightarrow V(Y_L)$ also holds in situations (a)-(b). Note that this concludes the proof of Theorem 1.10 since conjecture $V(X_L)$ (resp. $V(Y_L)$) holds when $\dim(X_L) \leq 2$ (resp. $\dim(Y_L) \leq 2$).

Let us now focus on class (i) of Theorem 1.7. The smooth projective k -schemes X_L and Y_L are of dimensions $8 - r$ and $r - 2$, respectively. Hence, $V(Y_L)$ holds when $r \leq 4$ and $V(X_L)$ when $r = 6$. When $r = 5$, X_L (and Y_L) is a Fano 3-fold. As explained by Gorchinskiy and Guletskii in [12, §5], the Chow motive of X admits a decomposition into Lefschetz motives and submotives of curves. This implies that $V(X_L)$ also holds.

Let us now focus on class (ii) of Theorem 1.7. The smooth projective k -schemes X_L and Y_L are of dimensions $10 - r$ and $r - 4$, respectively. Hence, $V(Y_L)$ holds when $r \leq 6$ and $V(X_L)$ when $r \geq 8$. This achieves the proof.

9. PROOF OF THEOREM 1.12

Following Kuznetsov [23, §4], let us denote by \mathfrak{a}_i the orthogonal complement of \mathbb{A}_{i+1} in \mathbb{A}_i ; these are called the ‘‘primitive subcategories’’ in *loc. cit.* Since the conjecture $V_{NC}(\mathbb{A}_0^{\text{dg}})$ holds, we hence conclude by induction that the conjectures $V_{NC}(\mathbb{A}_i^{\text{dg}})$ and $V_{NC}(\mathfrak{a}_i^{\text{dg}})$ also hold for every i . Thanks to HPD (see [23, Thm. 6.3]), $\text{perf}(Y)$ admits a Lefschetz decomposition $\text{perf}(Y) = \langle \mathbb{B}_m(-m), \dots, \mathbb{B}_0 \rangle$ with respect to $\mathcal{O}_Y(1)$ such that the primitive subcategories \mathfrak{b}_i coincide (via a Fourier-Mukai functor) with the primitive subcategories \mathfrak{a}_i . Consequently, $V_{NC}(\mathfrak{b}_i^{\text{dg}})$ holds for every i . An inductive argument, starting with $\mathfrak{b}_m = \mathbb{B}_m$, allows us then to conclude that conjecture $V_{NC}(\mathbb{B}_i^{\text{dg}})$ also holds for any i . Now, thanks once also to HPD (see [23, Thm. 5.3]), there exists also a triangulated category \mathbb{C}_L and semi-orthogonal decompositions

$$\begin{aligned} \text{perf}(X_L) &= \langle \mathbb{C}_L, \mathbb{A}_{\dim(L)}(1), \dots, \mathbb{A}_n(n - \dim(L)) \rangle \\ \text{perf}(Y_L) &= \langle \mathbb{B}_m(\dim(L^\perp) - m), \dots, \mathbb{B}_{\dim(L^\perp)}(-1), \mathbb{C}_L \rangle. \end{aligned}$$

Moreover, the composed functor $\text{perf}(X_L) \rightarrow \mathbb{C}_L \rightarrow \text{perf}(Y_L)$ is of Fourier-Mukai type. As a conclusion, since X_L and Y_L are smooth, we can apply Theorem 1.1 and obtain the following chain of equivalences

$$V(X_L) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(X_L)) \Leftrightarrow V_{NC}(\mathbb{C}_L^{\text{dg}}) \Leftrightarrow V_{NC}(\text{perf}_{\text{dg}}(Y_L)) \Leftrightarrow V(Y_L).$$

This achieves the proof.

10. PROOF OF THEOREM 1.14

Thanks to the work of Cossec [11], the conic bundle $Y \rightarrow \mathbb{P}^2$ has a natural structure of quartic double solid $Y \rightarrow \mathbb{P}^3$ ramified along a quartic symmetroid D . Via the natural involution on the resolution of singularities of D , one hence obtains an Enriques surface S ; consult [11, §3] for details. As proved by Zube in [38, §5], one has moreover a semi-orthogonal decomposition

$$\mathrm{perf}(S) = \langle \mathcal{T}_S, \mathcal{E}_1, \dots, \mathcal{E}_{10} \rangle,$$

with the \mathcal{E}_i 's exceptional objects. Let us denote by $\mathcal{T}_S^{\mathrm{dg}}$ the dg enhancement of \mathcal{T}_S induced from $\mathrm{perf}_{\mathrm{dg}}(S)$. Thanks to Lemma 2.1, $\mathcal{T}_S^{\mathrm{dg}}$ is smooth and proper. Hence, since S is a surface, an argument similar to the one of the proof of Theorem 1.2(i) shows us that conjecture $V_{NC}(\mathcal{T}_S^{\mathrm{dg}})$ holds.

Now, recall from Ingalls and Kuznetsov [14, §5.5] the construction of the Fourier-Mukai functor $\Phi : \mathrm{perf}(S) \rightarrow \mathrm{perf}(X)$ whose restriction to \mathcal{T}_S is fully-faithful. As proved in [14, Prop. 3.8 and Thm. 4.3], one has a semi-orthogonal decomposition

$$\mathrm{perf}(X) = \langle \Phi(\mathcal{T}_S), \mathcal{E}'_1, \mathcal{E}'_2 \rangle,$$

with the \mathcal{E}'_i 's exceptional objects. As a consequence, we obtain the equivalence

$$(10.1) \quad V_{NC}(\mathrm{perf}_{\mathrm{dg}}(X)) \Leftrightarrow V_{NC}(\Phi(\mathcal{T}_S)^{\mathrm{dg}}),$$

where $\Phi(\mathcal{T}_S)^{\mathrm{dg}}$ stands for the dg enhancement of $\Phi(\mathcal{T}_S)$ induced from $\mathrm{perf}_{\mathrm{dg}}(X)$. Since the kernel \mathcal{K} of the above Fourier-Mukai functor Φ gives rise to a Morita equivalence $\Phi_{\mathrm{dg}}^{\mathcal{K}} : \mathcal{T}_S^{\mathrm{dg}} \rightarrow \Phi(\mathcal{T}_S)^{\mathrm{dg}}$, we conclude that the conjecture (10.1) also holds. This achieves the proof.

Acknowledgments: The authors are grateful to Bruno Kahn and Claire Voisin for useful comments and answers. G. Tabuada was partially supported by a NSF CAREER Award.

REFERENCES

- [1] Y. André, *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*. Panoramas et Synthèses **17**. Société Mathématique de France, Paris, 2004.
- [2] Y. André and B. Kahn, *Nilpotence, radicaux et structures monoïdales*. (French) Rend. Sem. Mat. Univ. Padova **108** (2002), 107–291.
- [3] M. Artin, *Algebraization of formal moduli, II. Existence of modification*. Ann. of Math. **91**(2) (1970), 88–135
- [4] M. Artin and D. Mumford, *Some elementary examples of unirational varieties which are not rational*. Proc. London Math. Soc. (3) **25** (1972), 75–95.
- [5] A. Auel, M. Bernardara and M. Bolognesi, *Fibrations in complete intersections of quadrics, Clifford algebras, derived categories, and rationality problems*. Available at arXiv:1109.6938. To appear in Journal de Math. Pures et Appliquées.
- [6] H. Bass, *Algebraic K-theory*. W.A. Benjamin, New York, 1968.
- [7] M. Bernardara, M. Bolognesi and D. Faenzi, *Homological projective duality for determinantal varieties*. Available at arXiv:1410.7803.
- [8] M. Bernardara and G. Tabuada, *Chow groups of intersections of quadrics via homological projective duality and (Jacobians of) noncommutative motives*. Available at ArXiv:1310.6020.
- [9] A. Bondal and D. Orlov, *Derived categories of coherent sheaves*. International Congress of Mathematicians (Beijing), Vol II, 47–56, Higher Ed. Press, Beijing, 2002.
- [10] D.-C. Cisinski and G. Tabuada, *Symmetric monoidal structure on noncommutative motives*. Journal of K-Theory, **9** (2012), no. 2, 201–268.
- [11] F. Cossec, *Reye congruences*. Trans. AMS, Vol. **280** no. 2 (1983), p. 737–751.

- [12] S. Gorchinskiy and V. Guletskii, *Motives and representability of algebraic cycles on threefolds over a field* J. Algebraic Geom. **21** (2012), no. 2, 347–373.
- [13] T. Graber, J. Harris and J. Starr, *Families of rationally connected varieties*. J. Amer. Math. Soc. **16** (2003), no. 1, 57–67.
- [14] C. Ingalls and A. Kuznetsov, *On nodal Enriques surfaces and quartic double solids*. Available at arXiv:1012.3530.
- [15] B. Kahn and R. Sebastian, *Smash-nilpotent cycles on abelian 3-folds*. Math. Res. Lett. **16** (2009), no. 6, 1007–1010.
- [16] B. Keller, *On differential graded categories*. International Congress of Mathematicians (Madrid), Vol. II, 151–190, Eur. Math. Soc., Zürich, 2006.
- [17] M. Kontsevich, *Noncommutative motives*. Talk at the Institute for Advanced Study on the occasion of the 61st birthday of Pierre Deligne, October 2005. Video available at <http://video.ias.edu/Geometry-and-Arithmetic>.
- [18] ———, *Mixed noncommutative motives*. Talk at the Workshop on Homological Mirror Symmetry, Miami, 2010. Notes available at www-math.mit.edu/auroux/frg/miami10-notes.
- [19] ———, *Notes on motives in finite characteristic*. Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, 213–247, Progr. Math., **270**, Birkhuser Boston, MA, 2009.
- [20] A. Kuznetsov, *Base change for semiorthogonal decompositions*. Compositio Math. **147** (2011), no. 3, 852–876.
- [21] ———, *Derived categories of quadric fibrations and intersections of quadrics*. Advances in Math. **218** (2008), no. 5, 1340–1369.
- [22] ———, *Homological projective duality for Grassmannians of lines*. Available at arXiv:0610957.
- [23] ———, *Homological projective duality*, Publ. Math. Inst. Hautes Études Sci. (2007), no. 105, 157–220.
- [24] ———, *Semiorthogonal decompositions in algebraic geometry*. Available at arXiv:1404.3143
- [25] V. Lunts and D. Orlov, *Uniqueness of enhancement for triangulated categories*. J. Amer. Math. Soc. **23** (2010), 853–908.
- [26] M. Marcolli and G. Tabuada, *Noncommutative motives, numerical equivalence, and semi-simplicity*. American Journal of Mathematics **136** (2014), no. 1, 59–75.
- [27] ———, *Kontsevich’s noncommutative numerical motives*. Compositio Math. **148** (2012), no. 6, 1811–1820.
- [28] ———, *Noncommutative motives and their applications*. Available at arXiv:1311.2867.
- [29] ———, *Unconditional motivic Galois groups and Voevodsky’s nilpotence conjecture in the noncommutative world*. Available at arXiv:1112.5422(v1).
- [30] T. Matsusaka, *The criteria for algebraic equivalence and the torsion group*. Amer. J. Math. **79** (1957), 53–66.
- [31] B. G. Moishezon, *On n -dimensional compact varieties with n algebraically independent meromorphic functions, I, II and III*. Izv. Akad. Nauk SSSR Ser. Mat., **30**: 133–174.
- [32] G. Tabuada, *Chow motives versus noncommutative motives*. Journal of Noncommutative Geometry **7** (2013), no. 3, 767–786.
- [33] ———, *A guided tour through the garden of noncommutative motives*. Clay Mathematics Proceedings, Volume **16** (2012), 259–276.
- [34] G. Tabuada and M. Van den Bergh, *Noncommutative motives of Azumaya algebras*. Available at arXiv:1307.7946. To appear in Journal of the Institute of Mathematics of Jussieu.
- [35] C. Vial, *Algebraic cycles and fibrations*. Doc. Math. **18** (2013), 1521–1553.
- [36] V. Voevodsky, *A nilpotence theorem for cycles algebraically equivalent to zero*. Int. Math. Res. Notices (1995), no. 4, 187–198.
- [37] C. Voisin, *Remarks on zero-cycles of self-products of varieties*. Moduli of vector bundles (Sanda, 1994; Kyoto, 1994), 265–285, Lecture Notes in Pure and Appl. Math. **179**, Dekker, New York, 1996.
- [38] S. Zube, *Exceptional vector bundles on Enriques surfaces*. Mathematical Notes **61** (1997), no. 6, 693–699.

MARCELLO BERNARDARA, INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, 31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: marcello.bernardara@math.univ-toulouse.fr

URL: <http://www.math.univ-toulouse.fr/~mbernard/>

MATILDE MARCOLLI, MATHEMATICS DEPARTMENT, MAIL CODE 253-37, CALTECH, 1200 E. CALIFORNIA BLVD. PASADENA, CA 91125, USA

E-mail address: matilde@caltech.edu

URL: <http://www.its.caltech.edu/~matilde>

GONÇALO TABUADA, DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139, USA

E-mail address: tabuada@math.mit.edu

URL: <http://math.mit.edu/~tabuada>