

Quantum statistical mechanics, L -series, Anabelian Geometry

Matilde Marcolli

Beijing, August 2013

joint work with Gunther Cornelissen

General philosophy:

- Zeta functions are counting devices: spectra of operators with spectral multiplicities, counting ideals with given norm, number of periodic orbits, rational points, etc.
- Zeta function does not determine object: isospectral manifolds, arithmetically equivalent number fields, isogeny
- but ... sometimes a **family** of zeta functions does
- Zeta functions occur as partition functions of physical systems

Number fields: finite extensions \mathbb{K} of the field of rational numbers \mathbb{Q} .

- zeta functions: Dedekind $\zeta_{\mathbb{K}}(s)$ (for \mathbb{Q} Riemann zeta)
- symmetries: $G_{\mathbb{K}} = \text{Gal}(\overline{\mathbb{K}}/\mathbb{K})$ absolute Galois group; abelianized $G_{\mathbb{K}}^{ab}$
- adeles $\mathbb{A}_{\mathbb{K}}$ and ideles $\mathbb{A}_{\mathbb{K}}^*$, Artin map $\vartheta_{\mathbb{K}} : \mathbb{A}_{\mathbb{K}}^* \rightarrow G_{\mathbb{K}}^{ab}$
- topology: analogies with 3-manifolds (arithmetic topology)

How well do we understand them?

Analogy with manifolds: are there **complete invariants**?

Recovering a Number Field from invariants

- Dedekind zeta function $\zeta_{\mathbb{K}}(s) = \zeta_{\mathbb{L}}(s)$ **arithmetic equivalence**
Gaßmann examples:

$$\mathbb{K} = \mathbb{Q}(\sqrt[8]{3}) \text{ and } \mathbb{L} = \mathbb{Q}(\sqrt[8]{3 \cdot 2^4})$$

not isomorphism $\mathbb{K} \neq \mathbb{L}$

- Adeles rings $\mathbb{A}_{\mathbb{K}} \cong \mathbb{A}_{\mathbb{L}}$ **adelic equivalence** \Rightarrow arithmetic equivalence; Komatsu examples:

$$\mathbb{K} = \mathbb{Q}(\sqrt[8]{2 \cdot 9}) \text{ and } \mathbb{L} = \mathbb{Q}(\sqrt[8]{2^5 \cdot 9})$$

not isomorphism $\mathbb{K} \neq \mathbb{L}$

- Abelianized Galois groups: $G_{\mathbb{K}}^{\text{ab}} \cong G_{\mathbb{L}}^{\text{ab}}$ also not isomorphism;
Onabe examples:

$$\mathbb{K} = \mathbb{Q}(\sqrt{-2}) \text{ and } \mathbb{L} = \mathbb{Q}(\sqrt{-3})$$

not isomorphism $\mathbb{K} \neq \mathbb{L}$

- But ... absolute Galois groups $G_{\mathbb{K}} \cong G_{\mathbb{L}} \Rightarrow$ isomorphism
 $\mathbb{K} \cong \mathbb{L}$: Neukirch–Uchida theorem
(Grothendieck's **anabelian geometry**)

Question: Can combine $\zeta_{\mathbb{K}}(s)$, $\mathbb{A}_{\mathbb{K}}$ and $G_{\mathbb{K}}^{\text{ab}}$ to something as strong as $G_{\mathbb{K}}$ that determines isomorphism class of \mathbb{K} ?

Answer: Yes! Combine as a **Quantum Statistical Mechanical system**

Main Idea:

- Construct a QSM system associated to a number field
- Time evolution and equilibrium states at various temperatures
- Low temperature states are related to L-series
- Extremal equilibrium states determine the system
- System recovers the number field up to isomorphism

Purely number theoretic consequence:

An identity of all L -functions with Größencharakter gives an isomorphism of number fields

Quantum Statistical Mechanics (minimalist sketch)

- \mathcal{A} unital C^* -algebra of observables
- σ_t time evolution, $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{A})$
- states $\omega : \mathcal{A} \rightarrow \mathbb{C}$ continuous, normalized $\omega(1) = 1$, positive

$$\omega(a^*a) \geq 0$$

- equilibrium states $\omega(\sigma_t(a)) = \omega(a)$ all $t \in \mathbb{R}$
- representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$, Hamiltonian H


$$\pi(\sigma_t(a)) = e^{itH} \pi(a) e^{-itH}$$

- partition function $Z(\beta) = \text{Tr}(e^{-\beta H})$
- Gibbs states (equilibrium, inverse temperature β):

$$\omega_\beta(a) = \frac{\text{Tr}(\pi(a)e^{-\beta H})}{\text{Tr}(e^{-\beta H})}$$

- Generalization of Gibbs states: **KMS states**
 (Kubo–Martin–Schwinger) $\forall a, b \in A, \exists$ holomorphic $F_{a,b}$ on strip $I_\beta = \{0 < \text{Im } z < \beta\}$, bounded continuous on ∂I_β ,

$$F_{a,b}(t) = \omega(a\sigma_t(b)) \quad \text{and} \quad F_{a,b}(t + i\beta) = \omega(\sigma_t(b)a)$$



- Fixed $\beta > 0$: KMS_β state convex simplex: extremal states (like points in NCG)

Isomorphism of QSM systems: $\varphi : (\mathcal{A}, \sigma) \rightarrow (\mathcal{B}, \tau)$

$$\varphi : \mathcal{A} \xrightarrow{\cong} \mathcal{B}, \quad \varphi \circ \sigma = \tau \circ \varphi$$

C^* -algebra isomorphism intertwining time evolution

- Algebraic subalgebras $\mathcal{A}^\dagger \subset \mathcal{A}$ and $\mathcal{B}^\dagger \subset \mathcal{B}$: stronger condition: QSM isomorphism also preserves “algebraic structure”

$$\varphi : \mathcal{A}^\dagger \xrightarrow{\cong} \mathcal{B}^\dagger$$

- Pullback of a state: $\varphi^* \omega(a) = \omega(\varphi(a))$

Why QSM and Number theory? (a historical note)

1995: Bost–Connes QSM system $\mathcal{A}_{BC} = C(\hat{\mathbb{Z}}) \rtimes \mathbb{N}$

- generators $e(r)$, $r \in \mathbb{Q}/\mathbb{Z}$ and μ_n , $n \in \mathbb{N}$ and relations

$$\mu_n \mu_m = \mu_m \mu_n, \quad \mu_m^* \mu_m = 1$$

$$\mu_n \mu_m^* = \mu_m^* \mu_n \quad \text{if } (n, m) = 1$$

$$e(r + s) = e(r)e(s), \quad e(0) = 1$$

$$\mu_n e(r) \mu_n^* = \frac{1}{n} \sum_{ns=r} e(s)$$

- time evolution $\sigma_t(f) = f$ and $\sigma_t(\mu_n) = n^{it} \mu_n$

- representations $\pi_\rho : \mathcal{A}_{BC} \rightarrow \ell^2(\mathbb{N})$, $\rho \in \hat{\mathbb{Z}}^*$

$$\pi_\rho(\mu_n)\epsilon_m = \epsilon_{nm}, \quad \pi_\rho(\mathbf{e}(r))\epsilon_m = \zeta_r^m \epsilon_m$$

$\zeta_r = \rho(\mathbf{e}(r))$ root of unity

- Hamiltonian $H\epsilon_m = \log(m)\epsilon_m$, partition function

$$Z(\beta) = \text{Tr}(e^{-\beta H}) = \zeta_{\mathbb{Q}}(\beta)$$

Riemann zeta function

- Low temperature KMS states: L-series normalized by zeta
- Galois action on zero temperature states (class field theory)

Further generalizations: other QSM's with similar properties

- Bost-Connes as GL_1 -case of QSM for moduli spaces of \mathbb{Q} -lattices up to commensurability (Connes-M.M. 2006)
 $\Rightarrow GL_2$ -case, modular curves and modular functions
- QSM systems for imaginary quadratic fields (class field theory):
Connes-M.M.-Ramachandran
- B.Jacob and Consani-M.M.: QSM systems for function fields
(Weil and Goss L-functions as partition functions)
- Ha-Paugam: QSM systems for Shimura varieties \Rightarrow QSM
systems for arbitrary number fields (Dedekind zeta function)
further studied by Laca-Larsen-Neshveyev

We use these QSM systems for number fields

The **Noncommutative Geometry** viewpoint:

- Equivalence relation \mathcal{R} on X : quotient $Y = X/\mathcal{R}$. Even for very good $X \Rightarrow X/\mathcal{R}$ pathological!
- Functions on the quotient $\mathcal{A}(Y) := \{f \in \mathcal{A}(X) \mid f\mathcal{R} - \text{invariant}\}$
 \Rightarrow often too few functions: $\mathcal{A}(Y) = \mathbb{C}$ only constants
- NCG: $\mathcal{A}(Y)$ noncommutative algebra $\mathcal{A}(Y) := \mathcal{A}(\Gamma_{\mathcal{R}})$ functions on the graph $\Gamma_{\mathcal{R}} \subset X \times X$ of the equivalence relation with involution $f^*(x, y) = f(y, x)$ and convolution product

$$(f_1 * f_2)(x, y) = \sum_{x \sim u \sim y} f_1(x, u) f_2(u, y)$$

- $\mathcal{A}(\Gamma_{\mathcal{R}})$ associative noncommutative $\Rightarrow Y = X/\mathcal{R}$
noncommutative space (as good as X to do geometry, but new phenomena: time evolutions, thermodynamics, quantum phenomena)

In the various cases QSM system semigroup action on a space:

Bost–Connes revisited (Connes–M.M. 2006)

- \mathbb{Q} -lattices: (Λ, ϕ) \mathbb{Q} -lattice in \mathbb{R}^n : lattice $\Lambda \subset \mathbb{R}^n$ + group homomorphism

$$\phi : \mathbb{Q}^n / \mathbb{Z}^n \longrightarrow \mathbb{Q}\Lambda / \Lambda$$

- Commensurability: $(\Lambda_1, \phi_1) \sim (\Lambda_2, \phi_2)$ iff $\mathbb{Q}\Lambda_1 = \mathbb{Q}\Lambda_2$ and $\phi_1 = \phi_2 \pmod{\Lambda_1 + \Lambda_2}$
- Quotient \mathbb{Q} -lattices/Commensurability \Rightarrow NC space
- 1-dimensional \mathbb{Q} -lattices up to scaling $C(\hat{\mathbb{Z}})$

$$(\Lambda, \phi) = (\lambda \mathbb{Z}, \lambda \rho) \quad \lambda > 0$$

$$\rho \in \text{Hom}(\mathbb{Q}/\mathbb{Z}, \mathbb{Q}/\mathbb{Z}) = \varprojlim_n \mathbb{Z}/n\mathbb{Z} = \hat{\mathbb{Z}}$$

- with action of semigroup \mathbb{N} commensurability

$$\alpha_n(f)(\rho) = f(n^{-1}\rho) \quad \text{or zero}$$

$C(\hat{\mathbb{Z}}) \rtimes \mathbb{N}$ Bost–Connes algebra: moduli space

QSM systems for number fields: algebra and time evolution (A, σ)

$$A_{\mathbb{K}} := C(X_{\mathbb{K}}) \rtimes J_{\mathbb{K}}^+, \quad \text{with} \quad X_{\mathbb{K}} := G_{\mathbb{K}}^{\text{ab}} \times_{\hat{\sigma}_{\mathbb{K}}^*} \hat{\mathcal{O}}_{\mathbb{K}},$$

$\hat{\mathcal{O}}_{\mathbb{K}}$ = ring of finite integral adeles, $J_{\mathbb{K}}^+$ = is the semigroup of ideals, acting on $X_{\mathbb{K}}$ by Artin reciprocity

- Crossed product algebra $A_{\mathbb{K}} := C(X_{\mathbb{K}}) \rtimes J_{\mathbb{K}}^+$, generators and relations: $f \in C(X_{\mathbb{K}})$ and $\mu_{\mathfrak{n}}, \mathfrak{n} \in J_{\mathbb{K}}^+$

$$\mu_{\mathfrak{n}} \mu_{\mathfrak{n}}^* = e_{\mathfrak{n}}; \quad \mu_{\mathfrak{n}}^* \mu_{\mathfrak{n}} = 1; \quad \rho_{\mathfrak{n}}(f) = \mu_{\mathfrak{n}} f \mu_{\mathfrak{n}}^*;$$

$$\sigma_{\mathfrak{n}}(f) e_{\mathfrak{n}} = \mu_{\mathfrak{n}}^* f \mu_{\mathfrak{n}}; \quad \sigma_{\mathfrak{n}}(\rho_{\mathfrak{n}}(f)) = f; \quad \rho_{\mathfrak{n}}(\sigma_{\mathfrak{n}}(f)) = f e_{\mathfrak{n}}$$

- Artin reciprocity map $\vartheta_{\mathbb{K}} : \mathbb{A}_{\mathbb{K}}^* \rightarrow \mathbf{G}_{\mathbb{K}}^{\text{ab}}$, write $\vartheta_{\mathbb{K}}(\mathfrak{n})$ for ideal \mathfrak{n} seen as idele by non-canonical section s of

$$\mathbb{A}_{\mathbb{K},f}^* \begin{array}{c} \xrightarrow{\quad} \\ \searrow s \\ \xrightarrow{\quad} \end{array} \mathbf{J}_{\mathbb{K}} \quad : \quad (x_p)_p \mapsto \prod_{p \text{ finite}} p^{v_p(x_p)}$$

- semigroup action: $\mathfrak{n} \in \mathbf{J}_{\mathbb{K}}^+$ acting on $f \in C(X_{\mathbb{K}})$ as

$$\rho_{\mathfrak{n}}(f)(\gamma, \rho) = f(\vartheta_{\mathbb{K}}(\mathfrak{n})\gamma, s(\mathfrak{n})^{-1}\rho) e_{\mathfrak{n}},$$

$e_{\mathfrak{n}} = \mu_{\mathfrak{n}}\mu_{\mathfrak{n}}^*$ projector onto $[(\gamma, \rho)]$ with $s(\mathfrak{n})^{-1}\rho \in \hat{\mathcal{O}}_{\mathbb{K}}$

- partial inverse of semigroup action:

$$\sigma_{\mathfrak{n}}(f)(x) = f(\mathfrak{n} * x) \quad \text{with} \quad \mathfrak{n} * [(\gamma, \rho)] = [(\vartheta_{\mathbb{K}}(\mathfrak{n})^{-1}\gamma, \mathfrak{n}\rho)]$$

- Time evolution $\sigma_{\mathbb{K}}$ acts on $\mathbf{J}_{\mathbb{K}}^+$ as a phase factor $N(\mathfrak{n})^{it}$

$$\sigma_{\mathbb{K},t}(f) = f \quad \text{and} \quad \sigma_{\mathbb{K},t}(\mu_{\mathfrak{n}}) = N(\mathfrak{n})^{it} \mu_{\mathfrak{n}}$$

for $f \in C(\mathbf{G}_{\mathbb{K}}^{\text{ab}} \times \hat{\mathcal{O}}_{\mathbb{K}}^*)$ and for $\mathfrak{n} \in \mathbf{J}_{\mathbb{K}}^+$

Algebraic structure: **covariance algebra**

Algebraic subalgebra $A_{\mathbb{K}}^{\dagger}$ of C^* -algebra $A_{\mathbb{K}} := C(X_{\mathbb{K}}) \rtimes J_{\mathbb{K}}^+$:

$A_{\mathbb{K}}^{\dagger}$ unital, non-involutive algebra generated by $C(X_{\mathbb{K}})$ and the μ_n , $n \in J_{\mathbb{K}}^+$ (but not μ_n^*), with relations

$$\text{(using } \mu_n^* \mu_n = 1) \quad f \mu_n = \mu_n \sigma_n(f), \quad \mu_n f = \rho_n(f) \mu_n$$

Comment: presence of an algebraic subalgebra also in previous examples of arithmetic QSM

Comment: similar NCG interpretation as moduli spaces of \mathbb{K} -lattices up to commensurability

QSM isomorphism: two number fields \mathbb{K} and \mathbb{L}

$$\varphi : A_{\mathbb{K}} \xrightarrow{\sim} A_{\mathbb{L}}$$

C^* -algebra isomorphism

$$\varphi \circ \sigma_{\mathbb{K}} = \sigma_{\mathbb{L}} \circ \varphi$$

intertwines the time evolutions

$$\varphi : A_{\mathbb{K}}^{\dagger} \xrightarrow{\sim} A_{\mathbb{L}}^{\dagger}$$

preserves the covariance algebras

Theorem The following are equivalent:

- 1 $\mathbb{K} \cong \mathbb{L}$ are isomorphic number fields
- 2 Quantum Statistical Mechanical systems are isomorphic

$$(A_{\mathbb{K}}, \sigma_{\mathbb{K}}) \simeq (A_{\mathbb{L}}, \sigma_{\mathbb{L}})$$

C^* -algebra isomorphism $\varphi : A_{\mathbb{K}} \rightarrow A_{\mathbb{L}}$ compatible with time evolution, $\sigma_{\mathbb{L}} \circ \varphi = \varphi \circ \sigma_{\mathbb{K}}$ and covariance $\varphi : A_{\mathbb{K}}^{\dagger} \xrightarrow{\sim} A_{\mathbb{L}}^{\dagger}$

- 3 There is a group isomorphism $\psi : \hat{G}_{\mathbb{K}}^{ab} \rightarrow \hat{G}_{\mathbb{L}}^{ab}$ of Pontrjagin duals of abelianized Galois groups with

$$L_{\mathbb{K}}(\chi, s) = L_{\mathbb{L}}(\psi(\chi), s)$$

identity of all L -functions with Großencharakter

Comments:

- Generalization of arithmetic equivalence:
 $\chi = 1$ gives $\zeta_{\mathbb{K}}(s) = \zeta_{\mathbb{L}}(s)$
- Now also a purely number theoretic proof of (3) \Rightarrow (1) available by Hendrik Lenstra and Bart de Smit
- L -functions $L(\chi, s)$, for $s = \beta > 1$ is product of $\zeta_{\mathbb{K}}(\beta)$ and evaluation of an extremal KMS_{β} state of the QSM system $(A_{\mathbb{K}}, \sigma_{\mathbb{K}})$ at a test function $f_{\chi} \in C(X_{\mathbb{K}})$

Scheme of proof: (2) \Rightarrow (1)

- QSM isomorphism \Rightarrow arithmetic equivalence $\zeta_{\mathbb{K}}(s) = \zeta_{\mathbb{L}}(s)$
- $A_{\mathbb{K}}^{\dagger} \simeq A_{\mathbb{L}}^{\dagger}$ gives homeomorphism $X_{\mathbb{K}} \simeq X_{\mathbb{L}}$ and compatible semigroup isomorphism $J_{\mathbb{K}}^+ \simeq J_{\mathbb{L}}^+$
- Group isomorphism $G_{\mathbb{K}}^{ab} \simeq G_{\mathbb{L}}^{ab}$
- This preserves ramification \Rightarrow isomorphism of local units $\hat{\mathcal{O}}_{\wp}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\varphi(\wp)}^*$ and products $\varphi : \hat{\mathcal{O}}_{\mathbb{K}}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\mathbb{L}}^*$
- Semigroup isomorphism $A_{\mathbb{K},f}^* \cap \hat{\mathcal{O}}_{\mathbb{K}} \xrightarrow{\sim} A_{\mathbb{L},f}^* \cap \hat{\mathcal{O}}_{\mathbb{L}}$
- Endomorphism action of these \Rightarrow inner: $\mathcal{O}_{\mathbb{K},+}^{\times} \xrightarrow{\sim} \mathcal{O}_{\mathbb{L},+}^{\times}$
(tot pos non-zero integers)
- Recover additive structure (mod any totally split prime)
 $\varphi(x + y) = \varphi(x) + \varphi(y) \pmod{p}$

$$\Rightarrow \mathcal{O}_{\mathbb{K}} \simeq \mathcal{O}_{\mathbb{L}} \Rightarrow \mathbb{K} \simeq \mathbb{L}$$

Scheme of proof: (2) \Rightarrow (3)

- QSM isomorphism $\Rightarrow G_{\mathbb{K}}^{ab} \simeq G_{\mathbb{L}}^{ab}$ preserving ramification (as above)
- character groups $\psi : \hat{G}_{\mathbb{K}}^{ab} \xrightarrow{\sim} \hat{G}_{\mathbb{L}}^{ab}$
- character χ to function $f_{\chi} \in C(X_{\mathbb{K}})$, matching $\varphi(f_{\chi}) = f_{\psi(\chi)}$
- $\chi(\vartheta_{\mathbb{K}}(\mathbf{n})) = \psi(\chi)(\vartheta_{\mathbb{L}}(\varphi(\mathbf{n})))$
- Matching KMS $_{\beta}$ states: $\omega_{\gamma, \beta}^{\mathbb{L}}(\varphi(f)) = \omega_{\tilde{\gamma}, \beta}^{\mathbb{K}}(f)$
- using arithmetic equivalence: $L_{\mathbb{K}}(\chi, s) = L_{\mathbb{L}}(\psi(\chi), s)$

QSM isomorphism \Rightarrow **matching of L-series**

Scheme of proof: (3) \Rightarrow (1)

- need compatible isomorphisms $J_{\mathbb{K}}^+ \xrightarrow{\sim} J_{\mathbb{L}}^+$ and $C(X_{\mathbb{K}}) \xrightarrow{\sim} C(X_{\mathbb{L}})$
- know same number of primes \wp above same p with inertia degree f want to match compatibly with Artin map
- use combinations of L -series as counting functions: on finite quotients $\pi_G : G_{\mathbb{K}}^{ab} \rightarrow G$

$$\sum_{\substack{n \in J_{\mathbb{K}}^+ \\ N_{\mathbb{K}}(n)}} \left(\sum_{\widehat{G}} \chi(\pi_G(\gamma)^{-1}) \chi(\vartheta_{\mathbb{K}}(n)) \right) = b_{\mathbb{K}, G, n}(\gamma)$$

$$b_{\mathbb{K}, G, n}(\gamma) = \#\{n \in J_{\mathbb{K}}^+ : N_{\mathbb{K}}(n) = n \text{ and } \pi_G(\vartheta_{\mathbb{K}}(n)) = \pi_G(\gamma)\}$$

- For $G_{\mathbb{L}, n}^{ab} = \text{Gal}$ of max ab ext unram over n , get unique $m \in J_{\mathbb{L}}^+$ with $N_{\mathbb{L}}(m) = N_{\mathbb{K}}(n)$ and

$$\pi_{G_{\mathbb{K}, n}^{ab}}(\vartheta_{\mathbb{L}}(m)) = \pi_{G_{\mathbb{L}, n}^{ab}}((\psi^{-1})^*(\vartheta_{\mathbb{K}}(n)))$$

- Use stratification of $X_{\mathbb{K}}$ to extend $\psi : C(G_{\mathbb{K}}^{ab}) \xrightarrow{\sim} C(G_{\mathbb{L}}^{ab})$ to $\varphi : C(X_{\mathbb{K}}) \xrightarrow{\sim} C(X_{\mathbb{L}})$ compatibly with semigroup actions

One more equivalent formulation: \mathbb{K} and \mathbb{L} isomorphic iff \exists

- topological group isomorphism $\hat{\psi} : G_{\mathbb{K}}^{ab} \xrightarrow{\sim} G_{\mathbb{L}}^{ab}$
- semigroup isomorphism $\Psi : J_{\mathbb{K}}^+ \xrightarrow{\sim} J_{\mathbb{L}}^+$

with compatibility conditions

- Norm compatibility: $N_{\mathbb{L}}(\Psi(\mathfrak{n})) = N_{\mathbb{K}}(\mathfrak{n})$ for all $\mathfrak{n} \in J_{\mathbb{K}}^+$
- Artin map compatibility: for every finite abelian extension $\mathbb{K}' = (\mathbb{K}^{ab})^N / \mathbb{K}$, with $N \subset G_{\mathbb{K}}^{ab}$: prime \mathfrak{p} of \mathbb{K} unramified in \mathbb{K}'
 \Rightarrow prime $\Psi(\mathfrak{p})$ unramified in $\mathbb{L}' = (\mathbb{L}^{ab})^{\hat{\psi}(N)} / \mathbb{L}$ and

$$\hat{\psi}(\text{Frob}_{\mathfrak{p}}) = \text{Frob}_{\Psi(\mathfrak{p})}$$

Conclusions

- Is Quantum Statistical Mechanics a “noncommutative version” of anabelian geometry?
- What about function fields? QSM systems exist, purely NT proof seems not to work, but this QSM proof may work

General philosophy L -functions as coordinates determining underlying geometry

Examples:

- Cornelissen-M.M.: zeta functions of a spectral triple on limit set of Schottky uniformized Riemann surface determine conformal structure
- Cornelissen–J.W.de Jong: family of zeta functions of spectral triple of Riemannian manifold determine manifold up to isometry

Anabelian versus Noncommutative

- Anabelian geometry describes a number field \mathbb{K} in terms of the **absolute Galois group** $G_{\mathbb{K}}$
- But... no description of $G_{\mathbb{K}}$ in terms of **internal data** of \mathbb{K} only (Kronecker's hope)
- Langlands: relate to internal data via automorphic forms
- For **abelian** extensions yes: $G_{\mathbb{K}}^{ab}$ in terms of internal data: adeles, ideles (class field theory)
- But... $G_{\mathbb{K}}^{ab}$ does not recover \mathbb{K}
- Noncommutative geometry replaces $G_{\mathbb{K}}$ with the QSM system $(A_{\mathbb{K}}, \sigma_{\mathbb{K}})$ to reconstruct \mathbb{K}
- $A_{\mathbb{K}} = C(X_{\mathbb{K}}) \rtimes J_{\mathbb{K}}^+$ is built **only from internal data** of \mathbb{K} (primes, adeles, $G_{\mathbb{K}}^{ab}$)

More details on the proof of (2) \Rightarrow (1): **Stratification** of $X_{\mathbb{K}}$

- $\hat{\mathcal{O}}_{\mathbb{K},n} := \prod_{p|n} \hat{\mathcal{O}}_{\mathbb{K},p}$ and

$$X_{\mathbb{K},n} := G_{\mathbb{K}}^{\text{ab}} \times_{\hat{\mathcal{O}}_{\mathbb{K}}^*} \hat{\mathcal{O}}_{\mathbb{K},n} \quad \text{with} \quad X_{\mathbb{K}} = \varinjlim_n X_{\mathbb{K},n}$$

- Topological groups

$$G_{\mathbb{K}}^{\text{ab}} \times_{\hat{\mathcal{O}}_{\mathbb{K}}^*} \hat{\mathcal{O}}_{\mathbb{K},n}^* \simeq G_{\mathbb{K}}^{\text{ab}} / \vartheta_{\mathbb{K}}(\hat{\mathcal{O}}_{\mathbb{K},n}^*) = G_{\mathbb{K},n}^{\text{ab}}$$

Gal of max ab ext *unramified* at primes dividing n

- $J_{\mathbb{K},n}^+ \subset J_{\mathbb{K}}^+$ subsemigroup gen by prime ideals dividing n
- Decompose $X_{\mathbb{K},n} = X_{\mathbb{K},n}^1 \amalg X_{\mathbb{K},n}^2$

$$X_{\mathbb{K},n}^1 := \bigcup_{\mathfrak{n} \in J_{\mathbb{K},n}^+} \vartheta_{\mathbb{K}}(\mathfrak{n}) G_{\mathbb{K},n}^{\text{ab}} \quad \text{and} \quad X_{\mathbb{K},n}^2 := \bigcup_{p|n} Y_{\mathbb{K},p}$$

where $Y_{\mathbb{K},p} = \{(\gamma, \rho) \in X_{\mathbb{K},n} : \rho_p = 0\}$

- $X_{\mathbb{K},n}^1$ dense in $X_{\mathbb{K},n}$ and $X_{\mathbb{K},n}^2$ has $\mu_{\mathbb{K}}$ -measure zero
- Algebra $C(X_{\mathbb{K},n})$ is generated by functions

$$f_{\chi,n} : \gamma \mapsto \chi(\vartheta_{\mathbb{K}}(\mathfrak{n})) \chi(\gamma), \quad \chi \in \hat{G}_{\mathbb{K},n}^{\text{ab}}, \quad \mathfrak{n} \in J_{\mathbb{K},n}^+$$

First Step of (2) \Rightarrow (1): $(A_{\mathbb{K}}, \sigma_{\mathbb{K}}) \simeq (A_{\mathbb{L}}, \sigma_{\mathbb{L}}) \Rightarrow \zeta_{\mathbb{K}}(\mathbf{s}) = \zeta_{\mathbb{L}}(\mathbf{s})$

- QSM (A, σ) and representation $\pi : A \rightarrow B(\mathcal{H})$ gives Hamiltonian

$$\pi(\sigma_t(a)) = e^{itH} \pi(a) e^{-itH}$$

$$H_{\sigma_{\mathbb{K}}} \varepsilon_n = \log N(\mathbf{n}) \varepsilon_n$$

Partition function $\mathcal{H} = \ell^2(\mathcal{J}_{\mathbb{K}}^+)$

$$Z(\beta) = \text{Tr}(e^{-\beta H}) = \zeta_{\mathbb{K}}(\beta)$$

- Isomorphism $\varphi : (A_{\mathbb{K}}, \sigma_{\mathbb{K}}) \simeq (A_{\mathbb{L}}, \sigma_{\mathbb{L}}) \Rightarrow$ homeomorphism of sets of extremal KMS_{β} states by pullback $\omega \mapsto \varphi^*(\omega)$
- KMS_{β} states for $(A_{\mathbb{K}}, \sigma_{\mathbb{K}})$ classified [LLN]: $\beta > 1$

$$\omega_{\gamma, \beta}(f) = \frac{1}{\zeta_{\mathbb{K}}(\beta)} \sum_{\mathbf{m} \in \mathcal{J}_{\mathbb{K}}^+} \frac{f(\vartheta_{\mathbb{K}}(\mathbf{m})\gamma)}{N_{\mathbb{K}}(\mathbf{m})^{\beta}}$$

parameterized by $\gamma \in G_{\mathbb{K}}^{\text{ab}} / \vartheta_{\mathbb{K}}(\hat{\mathcal{O}}_{\mathbb{K}}^*)$

- Comparing GNS representations of $\omega \in \text{KMS}_\beta(\mathcal{A}_\mathbb{L}, \sigma_\mathbb{L})$ and $\varphi^*(\omega) \in \text{KMS}_\beta(\mathcal{A}_\mathbb{K}, \sigma_\mathbb{K})$ find Hamiltonians

$$H_\mathbb{K} = U H_\mathbb{L} U^* + \log \lambda$$

for some U unitary and $\lambda \in \mathbb{R}_+^*$

- Then partition functions give

$$\zeta_\mathbb{L}(\beta) = \lambda^{-\beta} \zeta_\mathbb{K}(\beta)$$

identity of Dirichlet series

$$\sum_{n \geq 1} \frac{a_n}{n^\beta} \quad \text{and} \quad \sum_{n \geq 1} \frac{b_n}{(\lambda n)^\beta}$$

with $a_1 = b_1 = 1$, taking limit as $\beta \rightarrow \infty$

$$a_1 = \lim_{\beta \rightarrow \infty} b_1 \lambda^{-\beta} \Rightarrow \lambda = 1$$

Conclusion of first step: **arithmetic equivalence** $\zeta_{\mathbb{L}}(\beta) = \zeta_{\mathbb{K}}(\beta)$

Consequences:

From arithmetic equivalence already know \mathbb{K} and \mathbb{L} have same degree over \mathbb{Q} , discriminant, normal closure, unit groups, archimedean places.

But... not class group (or class number)

Second Step of (2) \Rightarrow (1): unraveling the crossed product

$$\varphi : C(X_{\mathbb{K}}) \rtimes J_{\mathbb{K}}^+ \xrightarrow{\sim} C(X_{\mathbb{L}}) \rtimes J_{\mathbb{L}}^+ \quad \text{with} \quad \sigma_{\mathbb{L}} \circ \varphi = \varphi \circ \sigma_{\mathbb{K}}$$

and preserving the covariance algebra $\varphi : A_{\mathbb{K}}^{\dagger} \xrightarrow{\sim} A_{\mathbb{L}}^{\dagger}$

- Restrict to finitely many isometries μ_{\wp} , $N_{\mathbb{K}}(\wp) = p$
- $A_{\mathbb{K}}$ generated by $\mu_n f \mu_m^*$; in $A_{\mathbb{K}}^{\dagger}$ only $\mu_n f$
- Eigenspaces of time evolution in $A_{\mathbb{K}}^{\dagger}$ preserved:
so $C(X_{\mathbb{K}}) \xrightarrow{\sim} C(X_{\mathbb{L}})$ and $\varphi(\mu_n) = \sum \mu_m f_{n,m}$
- Commutators $[f, \mu_n] = (f - \rho_n(f))\mu_n$: match maximal ideals (mod commutators) so that homeomorphism $\Phi : X_{\mathbb{K}} \xrightarrow{\sim} X_{\mathbb{L}}$ compatible with semigroup actions $\gamma_{\alpha_x(n)}(\Phi(x)) = \Phi(\gamma_n(x))$ with locally constant $\alpha_x : J_{\mathbb{K}}^+ \rightarrow J_{\mathbb{L}}^+$ (that is, $\varphi(\mu_n) = \sum \delta_{m, \alpha_x(n)} \mu_m$)
- $\alpha_x = \alpha$ constant: know [LLN] ergodic action of $J_{\mathbb{K}}^+$ on $X_{\mathbb{K}}$, level sets would be clopen invariant subsets

Third Step of (2) \Rightarrow (1): isomorphism $G_{\mathbb{K}}^{ab} \xrightarrow{\sim} G_{\mathbb{L}}^{ab}$

- Projectors $e_{\mathbb{K},n} = \mu_n \mu_n^*$ mapped to projector $e_{\mathbb{L},\varphi(n)}$
- Fix $m \in J_{\mathbb{K}}^+$ and $\hat{\mathcal{O}}_{\mathbb{K},m} = \prod_{p|m} \hat{\mathcal{O}}_{\mathbb{K},p}$, then

$$V_{\mathbb{K},m} := \bigcap_{(m,n)=1} \text{Range}(e_{\mathbb{K},n}) = G_{\mathbb{K}}^{ab} \times_{\hat{\mathcal{O}}_{\mathbb{K}}^*} \{(0, \dots, 0, \hat{\mathcal{O}}_{\mathbb{K},m}, 0, \dots, 0)\}$$

$$\begin{aligned} \Phi(V_{\mathbb{K},m}) &= \bigcap_{(m,n)=1} \Phi(\text{Range}(e_{\mathbb{K},n})) = \bigcap_{(\varphi(m),\varphi(n))=1} \text{Range}(e_{\mathbb{L},\varphi(n)}) \\ &= G_{\mathbb{L}}^{ab} \times_{\hat{\mathcal{O}}_{\mathbb{L}}^*} \{(0, \dots, 0, \hat{\mathcal{O}}_{\mathbb{L},\varphi(m)}, 0, \dots, 0)\} = V_{\mathbb{L},\varphi(m)} \end{aligned}$$

- 1_m integral adèle = 1 at the prime divisors of m , zero elsewhere

$$H_{\mathbb{K},m} := G_{\mathbb{K}}^{ab} \times_{\hat{\mathcal{O}}_{\mathbb{K}}^*} \{1_m\} \subseteq X_{\mathbb{K}} \xrightarrow{\varphi} G_{\mathbb{L}}^{ab} \times_{\hat{\mathcal{O}}_{\mathbb{L}}^*} \{y_{\varphi(m)}\} \subseteq X_{\mathbb{L}}$$

- check that $y \in \hat{\mathcal{O}}_{\mathbb{L},m}^*$ is a unit
- then $H_{\mathbb{K},m}$ classes $[(\gamma, 1_m)] \sim [(\gamma', 1_m)] \iff \exists u \in \hat{\mathcal{O}}_{\mathbb{K}}^*$ with $\gamma' = \vartheta_{\mathbb{K}}(u)^{-1} \gamma$ and $1_m = u 1_m$

- then for $\mathring{G}_{\mathbb{K},m}^{ab}$ Gal of max ab ext unram *outside* prime div of m

$$H_{\mathbb{K},m} \cong G_{\mathbb{K}}^{ab} / \vartheta_{\mathbb{K}} \left(\prod_{q \nmid m} \hat{\sigma}_q^* \right) \cong \mathring{G}_{\mathbb{K},m}^{ab}$$

- $\mathring{G}_{\mathbb{K},m}^{ab}$ has dense subgroup gen by $\vartheta_{\mathbb{K}}(\mathfrak{n})$, ideals coprime to m
 $\Rightarrow H_{\mathbb{K},m}$ gen by these $\gamma_{\mathfrak{n}} := [(\vartheta_{\mathbb{K}}(\mathfrak{n})^{-1}, \mathbf{1}_m)]$
- with $\mathbf{1}_m = [(1, \mathbf{1}_m)]$ and $\Phi(\mathbf{1}_m) = [(x_m, y_m)]$ get

$$\Phi(\gamma_{\mathfrak{n}_1} \cdot \gamma_{\mathfrak{n}_2}) = \Phi([(\vartheta_{\mathbb{K}}(\mathfrak{n}_1 \mathfrak{n}_2)^{-1}, \mathbf{1}_m)])$$

$$= \Phi([(\vartheta_{\mathbb{K}}(\mathfrak{n}_1 \mathfrak{n}_2)^{-1}, \mathfrak{n}_1 \mathfrak{n}_2 \mathbf{1}_m)]) \text{ (since } \mathfrak{n}_1, \mathfrak{n}_2 \text{ coprime to } m)$$

$$= \Phi(\mathfrak{n}_1 \mathfrak{n}_2 * \mathbf{1}_m) = \varphi(\mathfrak{n}_1 \mathfrak{n}_2) * \Phi(\mathbf{1}_m) = [(\vartheta_{\mathbb{L}}(\varphi(\mathfrak{n}_1 \mathfrak{n}_2))^{-1} x_m, \varphi(\mathfrak{n}_1 \mathfrak{n}_2) y_m)]$$

- $\lim_{m \rightarrow +\infty} \mathbf{1}_m = 1 \Rightarrow \lim_{m \rightarrow +\infty} \Phi(\mathbf{1}_m) = \Phi(1)$ and get

$$\tilde{\Phi}(\gamma_1 \gamma_2) = \Phi(\gamma_1 \cdot \gamma_2) \Phi(1)^{-1} = \Phi(\gamma_1) \Phi(\gamma_2) \Phi(1)^{-2} = \tilde{\Phi}(\gamma_1) \cdot \tilde{\Phi}(\gamma_2)$$

Fourth step of (2): Preserving ramification

$N \subset G_{\mathbb{K}}^{\text{ab}}$ subgroup, $G_{\mathbb{K}}^{\text{ab}}/N \xrightarrow{\sim} G_{\mathbb{L}}^{\text{ab}}/\Phi(N)$

$$\mathfrak{p} \text{ ramifies in } \mathbb{K}'/\mathbb{K} \iff \varphi(\mathfrak{p}) \text{ ramifies in } \mathbb{L}'/\mathbb{L}$$

where $\mathbb{K}' = (\mathbb{K}^{\text{ab}})^N$ finite extension and $\mathbb{L}' := (\mathbb{L}^{\text{ab}})^{\Phi(N)}$

- seen have isomorphism $\Phi : \mathring{G}_{\mathbb{K}, \mathfrak{m}}^{\text{ab}} \xrightarrow{\sim} \mathring{G}_{\mathbb{L}, \varphi(\mathfrak{m})}^{\text{ab}}$ (Gal of max ab ext $\mathbb{K}_{\mathfrak{m}}$ unram outside prime div of \mathfrak{m})
- $\mathbb{K}' = (\mathbb{K}^{\text{ab}})^N$ fin ext ramified precisely above $\mathfrak{p}_1, \dots, \mathfrak{p}_r \in \mathcal{J}_{\mathbb{K}}^+$
- By previous $\mathbb{L}' := (\mathbb{L})^{\Phi(N)}$ contained in $\mathbb{L}_{\varphi(\mathfrak{p}_1) \dots \varphi(\mathfrak{p}_r)}$ but not in any $\mathbb{L}_{\varphi(\mathfrak{p}_1) \dots \widehat{\varphi(\mathfrak{p}_i)} \dots \varphi(\mathfrak{p}_r)}$ $\Rightarrow \mathbb{L}'/\mathbb{L}$ ramified precisely above $\varphi(\mathfrak{p}_1), \dots, \varphi(\mathfrak{p}_r)$

Fifth Step of (2) \Rightarrow (1): from QSM isomorphism get also

- Isomorphism of local units

$$\varphi : \hat{\mathcal{O}}_{\mathfrak{p}}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\varphi(\mathfrak{p})}^*$$

max ab ext where \mathfrak{p} unramified = fixed field of inertia group $I_{\mathfrak{p}}^{\text{ab}}$,
by ramification preserving

$$\Phi(I_{\mathfrak{p}}^{\text{ab}}) = I_{\varphi(\mathfrak{p})}^{\text{ab}}$$

and by local class field theory $I_{\mathfrak{p}}^{\text{ab}} \simeq \hat{\mathcal{O}}_{\mathfrak{p}}^*$

- by product of the local units: isomorphism

$$\varphi : \hat{\mathcal{O}}_{\mathbb{K}}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\mathbb{L}}^*$$

- Semigroup isomorphism

$$\varphi : (\mathbb{A}_{\mathbb{K},f}^* \cap \hat{\mathcal{O}}_{\mathbb{K}}, \times) \xrightarrow{\sim} (\mathbb{A}_{\mathbb{L},f}^* \cap \hat{\mathcal{O}}_{\mathbb{L}}, \times)$$

by exact sequence

$$0 \rightarrow \hat{\mathcal{O}}_{\mathbb{K}}^* \rightarrow \mathbb{A}_{\mathbb{K},f}^* \cap \hat{\mathcal{O}}_{\mathbb{K}} \rightarrow \mathcal{J}_{\mathbb{K}}^+ \rightarrow 0$$

(non-canonically) split by choice of uniformizer $\pi_{\mathfrak{p}}$ at every place

Recover **multiplicative** structure of the field

- **Endomorphism action** of $\mathbb{A}_{\mathbb{K},f}^* \cap \hat{\mathcal{O}}_{\mathbb{K}}$

$$\epsilon_s(f)(\gamma, \rho) = f(\gamma, s^{-1}\rho) e_{\tau}, \quad \epsilon_s(\mu_n) = \mu_n e_{\tau}$$

e_{τ} char function of set $s^{-1}\rho \in \hat{\mathcal{O}}_{\mathbb{K}}$

- $\hat{\mathcal{O}}_{\mathbb{K}}^*$ = part acting by automorphisms
- $\overline{\mathcal{O}_{\mathbb{K},+}^*}$ (closure of tot pos units): trivial endomorphisms
- $\mathcal{O}_{\mathbb{K},+}^{\times} = \mathcal{O}_{\mathbb{K},+} - \{0\}$ (non-zero tot pos elements of ring of integers): *inner endomorphisms* (isometries in $\mathbb{A}_{\mathbb{K}}^{\dagger}$ eigenv of time evolution)
- $\varphi(\epsilon_s) = \epsilon_{\varphi(s)}$ for all $s \in \mathbb{A}_{\mathbb{K},f}^* \cap \hat{\mathcal{O}}_{\mathbb{K}}$

Conclusion: isom of multiplicative semigroups of tot pos non-zero elements of rings of integers

$$\varphi : (\mathcal{O}_{\mathbb{K},+}^{\times}, \times) \xrightarrow{\sim} (\mathcal{O}_{\mathbb{L},+}^{\times}, \times)$$

Last Step of (2) \Rightarrow (1): Recover **additive** structure of the field

Extend by $\varphi(0) = 0$ the map $\varphi : (\mathcal{O}_{\mathbb{K},+}^{\times}, \times) \xrightarrow{\sim} (\mathcal{O}_{\mathbb{L},+}^{\times}, \times)$, Claim: it is additive

- Start with induced multipl map of local units $\varphi : \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\mathbb{L},\varphi(\mathfrak{p})}^*$ (from ramification preserving)
- set $\mathbf{1}_{\mathfrak{p}} = (0, \dots, 0, 1, 0, \dots, 0)$ and $\mathbf{1}_{\mathfrak{p}} := [(1, \mathbf{1}_{\mathfrak{p}})] \in X_{\mathbb{K}}$; for $u \in \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^*$, integral idele $u_{\mathfrak{p}} := (1, \dots, 1, u, 1, \dots, 1)$:
 $[(1, u_{\mathfrak{p}})] = [(\mathfrak{v}_{\mathbb{K}}(u_{\mathfrak{p}})^{-1}, 1)] \mapsto \Phi([(\mathfrak{v}_{\mathbb{K}}(u_{\mathfrak{p}})^{-1}, 1)]) =: [(1, \varphi(u))_{\varphi(\mathfrak{p})}]$
- Group isom to image $\lambda_{\mathbb{K},\mathfrak{p}} : \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* \rightarrow X_{\mathbb{K}} \xrightarrow{[\cdot \mathbf{1}_{\mathfrak{p}}]} Z_{\mathbb{K},\mathfrak{p}} \subset X_{\mathbb{K}}$
 $u \mapsto [(1, u_{\mathfrak{p}})] \mapsto [(1, u_{\mathfrak{p}} \cdot \mathbf{1}_{\mathfrak{p}})] = [(1, (0, \dots, 0, u, 0, \dots, 0))]$
- Commutative diagram

$$\begin{array}{ccc}
 \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* & \xrightarrow{\lambda_{\mathbb{K},\mathfrak{p}}} & Z_{\mathbb{K},\mathfrak{p}} \\
 \downarrow \varphi & & \downarrow \Phi \\
 \hat{\mathcal{O}}_{\mathbb{L},\varphi(\mathfrak{p})}^* & \xrightarrow{\lambda_{\mathbb{L},\varphi(\mathfrak{p})}} & Z_{\mathbb{L},\varphi(\mathfrak{p})}
 \end{array}$$

- Fix rational prime p totally split in \mathbb{K} (hence unramified) \Rightarrow arithm equiv: p tot split in \mathbb{L}
 - Set $\mathbb{Z}_{(p\Delta)}$ integers coprime to $p\Delta$ with $\Delta = \Delta_{\mathbb{K}} = \Delta_{\mathbb{L}}$ discriminant
 - map $\varpi_{\mathbb{K},p}: \mathbb{Z}_{(p\Delta)} \hookrightarrow \hat{\mathcal{O}}_{\mathbb{K},p}^* \rightarrow \mathbb{Z}_{\mathbb{K},p}$ with $\varpi_{\mathbb{K},p}: a \mapsto [(1, a \cdot \mathbf{1}_p)]$
 - $a = \mathfrak{p}_1 \dots \mathfrak{p}_r$ rational prime unramified \Rightarrow permute factors
 $\alpha_x((a)) = \mathfrak{p}_{\sigma(1)} \dots \mathfrak{p}_{\sigma(r)}$ so $\alpha_x((a)) = (a)$ fixes ideals $(a) \in \mathcal{J}_{\mathbb{Q}}^+$
- $$\Phi(\varpi_{\mathbb{K},p}(a)) = \Phi((a) * \mathbf{1}_p) = \alpha_{\mathbf{1}_p}((a)) * \Phi(\mathbf{1}_p) = (a) * \mathbf{1}_{\varphi(p)} = \varpi_{\mathbb{L},\varphi(p)}(a)$$
- so $\varphi: \hat{\mathcal{O}}_{\mathbb{K},p}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\mathbb{L},\varphi(p)}^*$ constant on $\mathbb{Z}_{(p\Delta)}$

- As above fix rational prime p totally split in \mathbb{K} (hence in \mathbb{L}) and $\mathfrak{p} \in \mathcal{J}_{\mathbb{K}}^+$ above p with $f(\mathfrak{p} | \mathbb{K}) = 1$ (hence $f(\varphi(\mathfrak{p}) | \mathbb{L}) = 1$)
- Use $\varphi : \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* \xrightarrow{\sim} \hat{\mathcal{O}}_{\mathbb{L},\varphi(\mathfrak{p})}^*$ to get multiplicative map of residue fields by Teichmüller lift $\tau_{\mathbb{K},p}: \overline{\mathbb{K}}_p^* \cong \mathbb{F}_p^* \hookrightarrow \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* \cong \mathbb{Q}_p^*$
- Show its extension by zero additive (hence identity map $\tilde{\varphi}: \mathbb{F}_p^* \rightarrow \mathbb{F}_p^*$) by extending $\tau_{\mathbb{K},p}: \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^* \rightarrow \hat{\mathcal{O}}_{\mathbb{K},\mathfrak{p}}^*$ with $x \mapsto \lim_{n \rightarrow +\infty} x^{p^n}$
- for \tilde{a} residue class in $\overline{\mathbb{K}}_p^* \cong \mathbb{F}_p$, choose integer a congruent to \tilde{a} mod p and coprime to discriminant Δ (Chinese remainder thm)

$$\varphi(\tau_{\mathbb{K},p}(a)) = \varphi \left(\lim_{n \rightarrow +\infty} a^{p^n} \right) = \lim_{n \rightarrow +\infty} \varphi(a)^{p^n} = \tau_{\mathbb{L},p}(\varphi(a)) = \tau_{\mathbb{L},p}(a)$$

$$\tilde{\varphi}(\tilde{a}) = \varphi(\tau_{\mathbb{K},p}(a)) \bmod \varphi(\mathfrak{p}) = \tau_{\mathbb{L},p}(a) \bmod \varphi(\mathfrak{p}) = \tilde{a} \bmod \varphi(\mathfrak{p})$$

- So φ identity mod any tot split prime, so for any $x, y \in \mathcal{O}_{\mathbb{K},+}$

$$\varphi(x + y) = \varphi(x) + \varphi(y) \bmod \varphi(\mathfrak{p})$$

- totally split primes of arbitrary large norm (Chebotarev)

$\Rightarrow \varphi$ **additive**