

Graph Hypersurfaces and a Dichotomy in the Grothendieck Ring

Letters in Mathematical Physics

A Journal for the Rapid
Dissemination of Short
Contributions in the Field of
Mathematical Physics

ISSN 0377-9017

Volume 95


Number 3

Lett Math Phys (2011)

95:223-232

DOI 10.1007/

s11005-011-0461-5



Your article is protected by copyright and all rights are held exclusively by Springer. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Graph Hypersurfaces and a Dichotomy in the Grothendieck Ring

PAOLO ALUFFI¹ and MATILDE MARCOLLI²¹*Department of Mathematics, Florida State University, Tallahassee, FL 32306, USA.**e-mail: aluffi@math.fsu.edu*²*Department of Mathematics, California Institute of Technology, Pasadena, CA 91125, USA. e-mail: matilde@caltech.edu*

Received: 1 June 2010 / Revised: 14 December 2010 / Accepted: 12 January 2011

Published online: 29 January 2011 – © Springer 2011

Abstract. The subring of the Grothendieck ring of varieties generated by the graph hypersurfaces of quantum field theory maps to the monoid ring of stable birational equivalence classes of varieties. We show that the image of this map is the copy of \mathbb{Z} generated by the class of a point. This clarifies the extent to which the graph hypersurfaces ‘generate the Grothendieck ring of varieties’: while it is known that graph hypersurfaces generate the Grothendieck ring over a localization of $\mathbb{Z}[\mathbb{L}]$ in which \mathbb{L} becomes invertible, the span of the graph hypersurfaces in the Grothendieck ring itself is nearly killed by setting the Lefschetz motive \mathbb{L} to zero. In particular, this shows that the graph hypersurfaces do *not* generate the Grothendieck ring prior to localization. The same result yields some information on the mixed Hodge structures of graph hypersurfaces, in the form of a constraint on the terms in their Deligne–Hodge polynomials. These observations are certainly not surprising for the expert reader, but are somewhat hidden in the literature. The treatment in this note is straightforward and self-contained.

Mathematics Subject Classification (2000). 81T15, 14E08, 16E20.

Keywords. graph hypersurfaces, Grothendieck ring, stable birational equivalence.

1. Introduction

The interplay between perturbative quantum field theory and the theory of motives of algebraic varieties has been extensively studied in recent years, in particular in terms of the algebro-geometric and motivic properties of the *graph hypersurfaces* associated with Feynman graphs of scalar quantum field theories.

In particular, one of the central results in the field is the main theorem of [6], which shows that graph hypersurfaces can be arbitrarily complicated from the motivic viewpoint: their affine complements are ‘*from the standpoint of their zeta functions, the most general schemes possible*’ ([6], p. 149). In rough terms, this is proven by showing that graph hypersurfaces generate the Grothendieck ring $K(\text{Var})$ of varieties.

The main purpose of this note is to clarify the extent to which this is the case, with emphasis on a subtlety which may be missed by a hurried reading of the literature. According to Theorem 0.6 in [6], graph hypersurfaces generate $S^{-1}K(\text{Var})$ as a module over the ring $S^{-1}\mathbb{Z}[\mathbb{L}]$, where S is the (saturated) multiplicative subset of $\mathbb{Z}[\mathbb{L}]$ generated by $\mathbb{L}^n - \mathbb{L}$ for $n > 1$, with $\mathbb{L} = [\mathbb{A}^1]$ the Lefschetz–Tate motive. This localization is in fact necessary to the result of [6], and we give a simple geometric explanation of why this is the case. We will point out that graph hypersurfaces do not generate the Grothendieck ring as a module over $\mathbb{Z}[\mathbb{L}]$, and in fact they do not even generate the localization $S'^{-1}K(\text{Var})$ as a module over $S'^{-1}\mathbb{Z}[\mathbb{L}]$, where S' is generated by $\mathbb{L}^n - 1$ for $n > 0$. For example, we will show that the class of an elliptic curve is not in the span of the graph hypersurfaces if coefficients are taken in $S'^{-1}\mathbb{Z}[\mathbb{L}]$. As $S^{-1}\mathbb{Z}[\mathbb{L}]$ is the localization of this latter ring at \mathbb{L} , it appears that localization at \mathbb{L} is crucial the result of [6]. Considering this observation, we propose the following

QUESTION . *Do graph hypersurfaces generate the localized Grothendieck ring as a module over $\mathbb{Z}[\mathbb{L}, \mathbb{L}^{-1}]$?*

This situation illustrates a sharp dichotomy in the behavior of graph hypersurfaces in the Grothendieck ring under the contrasting operations of inverting \mathbb{L} and setting \mathbb{L} to zero. Graph hypersurfaces are ‘as general as possible’ after localization (at \mathbb{L} and S'), while they are extremely special with respect to taking the quotient modulo the ideal (\mathbb{L}) : their span agrees with the span of a point modulo (\mathbb{L}) . While these facts are certainly not surprising to the expert, we feel that it is useful to lay them out in a short and self-contained note, with clearly stated formulas and transparent proofs. As we will point out at the end of this introduction, stronger statements can be proven concerning the triviality modulo (\mathbb{L}) , for physically significant graphs, but at the price of more combinatorially demanding arguments.

We quickly recall some basic notation and terminology. For a connected finite graph G with n edges, the graph polynomial $\psi_G(t_1, \dots, t_n)$ is defined as

$$\psi_G(t_1, \dots, t_n) = \sum_{T \subseteq G} \prod_{e \notin T} t_e,$$

where T runs over the spanning trees of G and t_e is the variable associated with an edge e . In general, we define the graph polynomial for a (finite) graph G to be the product of the polynomials for the connected components of G . We denote by X_G the projective hypersurface defined by the homogeneous polynomial ψ_G in \mathbb{P}^{n-1} , by $\widehat{X}_G \subset \mathbb{A}^n$ the affine hypersurface, and by $Y_G \subseteq \mathbb{A}^n$ the affine hypersurface complement $Y_G = \mathbb{A}^n \setminus \widehat{X}_G$. As the main results of [6] are expressed in terms of Y_G , we choose to focus on Y_G in this paper. We note that graph hypersurfaces are usually singular; also, it is easy to see that the irreducible components of X_G are rational.

Our main result can be stated as follows. Larsen and Lunts associate with each variety V (possibly singular, possibly non-compact) an element in the monoid ring $\mathbb{Z}[SB]$ generated by *stable birational equivalence* classes of varieties. This assignment is compatible with the relations defining $K(\text{Var})$, and associates with V its own stable birational equivalence class *if V is smooth and projective*. Smooth projective rational varieties have class 1 in $\mathbb{Z}[SB]$, but this is in general not the case for rational varieties that are singular or not complete: the element in $\mathbb{Z}[SB]$ determined by the image of the class $[V] \in K(\text{Var})$ of a quasi-projective or singular rational variety need not be in the ‘constant’ part $\mathbb{Z} \subseteq \mathbb{Z}[SB]$ (cf. Example 2.4). Thus, although irreducible graph hypersurfaces are rational, this fact alone does not give information on their image in $\mathbb{Z}[SB]$. What we show is precisely that the Larsen–Lunts images of graph hypersurfaces *do* lie in the constant part of $\mathbb{Z}[SB]$.

THEOREM 1.1. *Affine graph hypersurface complements span $\mathbb{Z} \subseteq \mathbb{Z}[SB]$.*

Morally, Theorem 1.1 shows that graph hypersurfaces and their complements are rational in a very strong sense (which we will make precise in Sect. 2): for example, the image in $\mathbb{Z}[SB]$ of the class of an irreducible graph hypersurface does equal the class of a point (Corollary 3.3). This is the reason why they do not span the unlocalized Grothendieck ring of varieties.

In more classical terms, Theorem 1.1 provides some information on the mixed Hodge structure of graph hypersurfaces, as it shows that the Deligne–Hodge polynomial of a (projective) graph hypersurface X_G is necessarily of the form $c + uvP(u, v)$, with $c \in \mathbb{Z}$. This shows that elliptic curves are not in the span of graph hypersurfaces in the Grothendieck ring.

Theorem 1.1 is proven using the realization of $\mathbb{Z}[SB]$ as the quotient $K(\text{Var})/(\mathbb{L})$ (also recalled in Section 2). An explicit computation (Theorem 3.2) based on a deletion–contraction formula for the Grothendieck class of a graph hypersurface shows that for every graph G , the class of the complement Y_G modulo (\mathbb{L}) is 0 or ± 1 . Theorem 1.1 and the consequences mentioned above follow immediately.

We remark that Theorem 3.2 is certainly not the strongest possible result, although it suffices for our purposes and is extremely easy to prove, as we show in this paper. Lemma 15 in [8] shows that for ‘physically significant’ graphs G the class of Y_G is in fact 0 modulo (\mathbb{L}^2) (see also Corollary 2.7 in [13]; this fact ultimately relies on Dodgson’s identity). For these graphs, the Deligne–Hodge polynomial of X_G has the form $1 + uv + u^2v^2P(u, v)$. ‘Banana graphs’ give examples of graphs which do not satisfy this stronger property: the class for the n -banana graph is a multiple of \mathbb{L} , but not of \mathbb{L}^2 for $n \neq 3$ ([2], Corollary 3.13).

2. Stable Birational Equivalence and the Grothendieck Ring

In the following, we denote by $K(\text{Var})$ the Grothendieck ring of varieties. This is generated by the isomorphism classes of irreducible quasi-projective varieties with

the inclusion–exclusion relations $[X]=[X \setminus Y]+[Y]$, for closed embeddings $Y \subseteq X$, and with the product $[X] \cdot [Y]=[X \times Y]$. The Grothendieck ring $K(\text{Var})$ depends on the field of definition of the varieties. This will be understood to be \mathbb{Q} in the following.

A result of [12] relates the Grothendieck ring to the ring of stable birational equivalence classes of varieties. We comment here briefly on some aspects of this result that will be useful in the discussion of the case of graph hypersurfaces.

Two (irreducible, complex) varieties X, Y are *stably birational* if $X \times \mathbb{P}^k$ is birational to $Y \times \mathbb{P}^\ell$ for some $k, \ell \geq 0$. If X is stably birational to Y , and X' is stably birational to Y' , then $X \times X'$ is stably birational to $Y \times Y'$. Thus, the set of classes of stable birational equivalence of varieties is a multiplicative monoid SB , with unit equal to the class of a point.

THEOREM 2.1. ([12], Proposition 2.7) *Let (\mathbb{L}) be the ideal in $K(\text{Var})$ generated by the Lefschetz motive $\mathbb{L}=[\mathbb{A}^1]$. The ring $K(\text{Var})/(\mathbb{L})$ is isomorphic to the monoid ring $\mathbb{Z}[SB]$.*

This result is obtained by defining a homomorphism $K(\text{Var}) \rightarrow \mathbb{Z}[SB]$, sending $[V]$ to the stable birational equivalence class $[V]_{SB}$ of V for every irreducible *smooth, projective* variety V . The main technical step is to show that this homomorphism is well defined; this may be proven by using Bittner's alternative description ([5]) of the Grothendieck ring of varieties $K(\text{Var})$ with generators that are smooth projective varieties and relations

$$[X] - [Y] = [Bl_Y(X)] - [E],$$

for a smooth closed subvariety $Y \subseteq X$, with $Bl_Y(X)$ the blowup of X along Y and E the exceptional divisor. This relation replaces the usual inclusion-exclusion relation $[X]=[X \setminus Y]+[Y]$, which requires the noncompact $X \setminus Y$. Bittner's characterization depends on the *weak factorization theorem* of [1], which shows that any proper birational map between smooth irreducible varieties over a field of characteristic zero can be factored into a sequence of blow-ups and blow-downs with smooth centers.

Theorem 2.1 is stated over \mathbb{C} in [12], but holds over \mathbb{Q} as well since so does the weak factorization theorem (Remark 2 after Theorem 0.3.1 of [1]); this is also observed explicitly in [11], p. 28.

Remark 2.2. Stable birational equivalence makes sense for every variety V , so every variety (whether or not smooth and projective) has a class $[V]_{SB}$ in $\mathbb{Z}[SB]$. It is important to keep in mind that in general this class agrees with the image of $[V] \in K(\text{Var})$ via the Larsen-Lunts homomorphism *only if* V is smooth and projective. In other cases the image of $[V]$ in $\mathbb{Z}[SB]$ may be determined by expressing

$[V]$ as a combination of classes of smooth projective varieties, and then reproducing that combination in $\mathbb{Z}[SB]$.

For example, the image of $\mathbb{L} = [\mathbb{A}^1]$ is 0 in $\mathbb{Z}[SB]$ because $[\mathbb{A}^1] = [\mathbb{P}^1] - [\mathbb{P}^0]$ in $K(\text{Var})$, and $\mathbb{P}^0, \mathbb{P}^1$ are trivially stably birationally equivalent to each other. Likewise, the image of (the class of) an irreducible nodal plane cubic C in $\mathbb{Z}[SB]$ is $0 \neq 1$ even though $[C]_{SB} = [\mathbb{P}^0]_{SB}$, since $[C] = [\mathbb{P}^1] - [\mathbb{P}^0]$ in $K(\text{Var})$.

To make this point more explicit, we introduce a notion of ‘ \mathbb{L} -equivalence’.

DEFINITION 2.3. Two irreducible quasi-projective varieties X, Y are \mathbb{L} -equivalent if their classes in $\mathbb{Z}[SB]$ via the Larsen–Lunts isomorphism coincide, that is, if $[X] \equiv [Y] \pmod{(\mathbb{L})}$ in $K(\text{Var})$. A variety is \mathbb{L} -rational if it is \mathbb{L} -equivalent to \mathbb{P}^k , for some $k \geq 0$.

If X and Y are irreducible smooth projective and stably birational, then they are also \mathbb{L} -equivalent; however, this is not necessarily the case if X, Y are not smooth and/or not complete. For example, as observed above, an irreducible nodal cubic in \mathbb{P}^2 is complete and birational to \mathbb{P}^1 but not \mathbb{L} -rational.

In fact, the following example shows that rational (singular, projective) varieties may be very far from being \mathbb{L} -rational.

EXAMPLE 2.4. There exists a complete rational surface X whose Larsen–Lunts image in $\mathbb{Z}[SB]$ is $2 - [C]_{SB}$, where C is an elliptic curve.

Indeed, by Theorem 3.3 of [10], there exist projective rational surfaces X with one isolated singular point p such that the exceptional divisor in the minimal resolution \tilde{X} of X is an elliptic curve C . Using Bittner’s relations, $[X] = [\tilde{X}] - [C] + [p]$ in $K(\text{Var})$, and since all varieties on the right hand side are smooth and projective, and \tilde{X} is rational, then the image of $[X]$ in $\mathbb{Z}[SB]$ equals $[\tilde{X}]_{SB} - [C]_{SB} + [p]_{SB} = 2 - [C]_{SB}$.

These caveats apply to graph hypersurfaces. As recalled in the introduction, irreducible (projective) graph hypersurfaces X_G are easily seen to be rational, and are complete, but are in general singular. Affine graph hypersurface complements Y_G are trivially rational, but non-complete. As observed above, these naive considerations do not suffice to determine the Larsen–Lunts images of these varieties in $\mathbb{Z}[SB]$.

3. Graph Hypersurfaces and Stable Birational Equivalence

In [4] we proved a deletion-contraction formula for the classes in the Grothendieck ring of the graph hypersurfaces. We recall here the result, giving a short and self-contained proof for completeness, since the conclusion we derive on the stable birational equivalence classes will be a direct consequence of this formula. The

result is also implicit in the literature preceding [4], e.g. [7,14]. However, the formula given for $[X_\Gamma]$ in [7] (8.8) appears to be incorrect as stated, as simple examples show (for instance, \Leftrightarrow). The isomorphism given in (8.2) of loc. cit. is correct, and does imply the formula given in the following theorem.

THEOREM 3.1. ([4], Theorem 3.8) *Let G be a graph with n edges, and let e be an edge of G . Denote by $G \setminus e$ the graph obtained by removing e , and by G/e the graph obtained by contracting e . Let Y_G denote the affine graph hypersurface complement in \mathbb{A}^n .*

- *If e is a bridge in G , then $[Y_G] = \mathbb{L} \cdot [Y_{G \setminus e}]$;*
- *If e is a looping edge in G , then $[Y_G] = (\mathbb{L} - 1) \cdot [Y_{G \setminus e}]$;*
- *If e is neither a bridge nor a looping edge in G , then*

$$[Y_G] = \mathbb{L} \cdot [\mathbb{A}^{n-1} \setminus Z] - [Y_{G \setminus e}],$$

where Z is the intersection of the affine graph hypersurfaces for $G \setminus e$, G/e in \mathbb{A}^{n-1} .

Proof. We give a quick proof for completeness; for more details, see [4]. Let t_i be the variable corresponding to the i -th edge e_i , and assume that $e = e_n$ is not a bridge or a looping edge. In terms of graph polynomials:

$$P_G(t_1, \dots, t_n) = t_n P_{G \setminus e}(t_1, \dots, t_{n-1}) + P_{G/e}(t_1, \dots, t_{n-1}).$$

The complement Y_G is the set of n -tuples (t_1, \dots, t_n) such that $P_G(t_1, \dots, t_n)$ does not vanish; thus, such that

$$t_n P_{G \setminus e}(t_1, \dots, t_{n-1}) \neq -P_{G/e}(t_1, \dots, t_{n-1}).$$

Over $Y_{G \setminus e}$ (that is, if $P_{G \setminus e} \neq 0$), this condition holds if and only if $t_n \neq -P_{G/e}/P_{G \setminus e}$, that is, for $t_n \in \mathbb{A}^1 \setminus \mathbb{A}^0$. Over the rest of the complement of Z (that is, if $P_{G \setminus e} \neq 0$ but $P_{G/e} = 0$), the condition is satisfied for all t_n , hence for $t_n \in \mathbb{A}^1$. Over Z (that is, if $P_{G \setminus e} = P_{G/e} = 0$), the condition is not satisfied for any choice of t_n . Therefore,

$$[Y_G] = (\mathbb{L} - 1) \cdot [Y_{G \setminus e}] + \mathbb{L} \cdot [(\mathbb{A}^{n-1} \setminus Z) \setminus Y_{G \setminus e}],$$

and this is equivalent to third equality stated above. The other cases are analogous. □

The deletion–contraction formula yields the following computation of the Larsen–Lunts image of Y_G in $\mathbb{Z}[SB] \cong K(\text{Var})/(\mathbb{L})$.

THEOREM 3.2.

$$[Y_G] \equiv \begin{cases} 0 & \text{mod } (\mathbb{L}) \text{ if } G \text{ has edges that are not looping edges} \\ (-1)^n & \text{mod } (\mathbb{L}) \text{ if } G \text{ has } n \geq 0 \text{ looping edges, and no other edge.} \end{cases}$$

Proof. Reading the result of Theorem 3.1 above modulo (\mathbb{L}) gives:

- If e is a bridge in G , then $[Y_G] = 0 \pmod{(\mathbb{L})}$;
- If e is not a bridge G , then $[Y_G] = -[Y_{G \setminus e}] \pmod{(\mathbb{L})}$.

If now G has any edge that is not a looping edge, then removing all but one such edge leaves a graph with a bridge, and hence $[Y_G] \equiv 0 \pmod{(\mathbb{L})}$ in this case. If all n edges of G are looping edges, then repeated application of the second formula shows that $[Y_G] \equiv (-1)^n \cdot [Y_{\underline{G}}] \pmod{(\mathbb{L})}$, where \underline{G} is the graph obtained by removing all edges from G . Clearly $Y_{\underline{G}} = \mathbb{A}^0$, so $[Y_G] \equiv (-1)^n \pmod{(\mathbb{L})}$ in this case, as stated. \square

In terms of projective graph hypersurfaces:

COROLLARY 3.3. *Let G be a graph that is not a forest, and with at least one non-looping edge. Then the projective graph hypersurface X_G is \mathbb{L} -rational.*

Proof. Note that G must have at least 2 edges. Since G is not a forest, \widehat{X}_G is not empty; and by Theorem 3.2, since G has non-looping edges, then the class $[Y_G]$ is in the ideal (\mathbb{L}) . The affine complement $Y_G = \mathbb{A}^n \setminus \widehat{X}_G$ fibers over the projective complement $\mathbb{P}^{n-1} \setminus X_G$, with fibers $\mathbb{A}^1 \setminus \mathbb{A}^0$. Therefore

$$(\mathbb{L} - 1) \cdot [\mathbb{P}^{n-1} \setminus X_G] \in (\mathbb{L}),$$

and hence $[\mathbb{P}^{n-1} \setminus X_G] \in (\mathbb{L})$; thus

$$[X_G] \equiv [\mathbb{P}^{n-1}] \pmod{(\mathbb{L})},$$

showing that X_G is \mathbb{L} -equivalent to \mathbb{P}^{n-1} . \square

Again we remark that if X_G is irreducible, then it is easily seen to be rational, but examples such as Example 2.4 show that this does not suffice in itself to draw the conclusion stated in Corollary 3.3.

Considering the result of [12] recalled in Theorem 2.1 above, Theorem 3.2 has the following immediate consequence:

COROLLARY 3.4. *Let \mathcal{R} be the subring of $K(\text{Var})$ spanned by the classes $[Y_G]$. Then the image of \mathcal{R} in $\mathbb{Z}[SB]$ via the Larsen–Lunts homomorphism is the subring \mathbb{Z} generated by the stable birational equivalence class of a point.*

Proof. The image of \mathcal{R} in $\mathbb{Z}[SB]$ is the quotient $\mathcal{R}/(\mathbb{L})$. By Theorem 3.2, $\mathcal{R}/(\mathbb{L}) \cong \mathbb{Z}$. \square

Remark 3.5. With our conventions, the product of two classes $[Y_{G_1}]$, $[Y_{G_2}]$ is itself the class $[Y_{G_1 \sqcup G_2}]$ of the affine complement of a graph hypersurface (the class of

the affine hypersurface complement is a ‘motivic Feynman rule’, see Proposition 2.5 in [3]). Further, the Lefschetz motive \mathbb{L} equals $[Y_G]$ for the graph G consisting of a single edge joining two distinct vertices. Thus, the ring \mathcal{R} generated by the classes $[Y_G]$ agrees with the $\mathbb{Z}[\mathbb{L}]$ -module generated by the classes $[Y_G]$. Therefore, the following immediate consequence of Corollary 3.4 formalizes the first ‘non-spanning’ result mentioned in the Introduction.

THEOREM 3.6. *The classes $[Y_G]$ of the affine graph hypersurface complements do not span the Grothendieck ring $K(\text{Var})$ over $\mathbb{Z}[\mathbb{L}]$.*

Proof. Considering Corollary 3.4, it suffices to notice that $\mathbb{Z}[SB] \neq \mathbb{Z}$. This is clear since there are polynomial invariants of smooth projective varieties that are invariant under stable birational equivalence, and are not always constant. One example is given in [12], Definition 3.4 in terms of Hodge polynomials. \square

As mentioned in the Introduction, this observation can be sharpened, in a way that relates well to the result of [6].

COROLLARY 3.7. *Let S' be the saturated multiplicative subset of $\mathbb{Z}[\mathbb{L}]$ generated by the elements $\mathbb{L}^n - 1$, for $n > 0$. Then the classes $[Y_G]$ do not generate $S'^{-1}K(\text{Var})$ over $S'^{-1}\mathbb{Z}[\mathbb{L}]$.*

Proof. Localization commutes with taking quotients: the quotient

$$(S'^{-1}K(\text{Var})) / (\mathbb{L})$$

equals the localization $S'^{-1}(K(\text{Var}) / (\mathbb{L}))$. Since all elements of S' are invertible modulo (\mathbb{L}) , this latter equals $K(\text{Var}) / (\mathbb{L})$; and the action of \mathbb{L} on this module is 0. Since the classes $[Y_G]$ do not span this quotient, they cannot span $S'^{-1}K(\text{Var})$ over $S'^{-1}\mathbb{Z}[\mathbb{L}]$. \square

Corollary 3.7 should be compared with Theorem 0.6 in [6], which states that the classes $[Y_G]$ do generate the localization $S^{-1}K(\text{Var}_{\mathbb{Z}})$ over $S^{-1}\mathbb{Z}[\mathbb{L}]$, where S is the multiplicative system generated by $\mathbb{L}^n - \mathbb{L}$ for $n > 1$. *A fortiori*, the classes $[Y_G]$ generate $S^{-1}K(\text{Var}_{\mathbb{Q}})$. Localizing at S amounts to localizing at S' and at \mathbb{L} ; Corollary 3.7 shows that the localization at \mathbb{L} is crucial to the mentioned result in [6]. This suggests that the classes $[Y_G]$ may possibly span the Grothendieck ring over the simpler localization $\mathbb{Z}[\mathbb{L}^{-1}, \mathbb{L}]$.

We end by observing that the results stated above have a straightforward Hodge-theoretic formulation. Every smooth complex projective variety X carries Hodge numbers $h^{p,q}(X)$. The *Hodge polynomial* of X is the polynomial $\sum_{p,q} (-1)^{p+q} h^{p,q}(X) u^p v^q$. Now, the Hodge polynomial determines a ring homomorphism $K(\text{Var}) \rightarrow \mathbb{Z}[u, v]$, see e.g. Section 2.11 in [15]; this homomorphism

maps \mathbb{L} to uv . Indeed, the Hodge polynomial may be consistently defined for all varieties and satisfies the relations in $K(\text{Var})$, as observed in [9]. This (‘Deligne–Hodge’) polynomial of an arbitrary complex variety X records information about the mixed Hodge structure on the cohomology $H_c^k(X, \mathbb{Q})$ of X with compact support: it may be defined as $\sum_{p,q} e^{p,q}(X)u^p v^q$, where

$$e^{p,q}(X) = \sum_k (-1)^k h^{p,q}(H_c^k(X, \mathbb{Q})).$$

COROLLARY 3.8. *Let X be any complex projective variety whose class $[X]$ is in the subring \mathcal{R} of $K(\text{Var}_{\mathbb{C}})$ generated by the classes $[Y_G]$, as G ranges over all graphs. Then the Deligne–Hodge polynomial of X is of the form*

$$c + uvP(u, v), \tag{3.1}$$

with $c \in \mathbb{Z}$.

Proof. The Hodge polynomial induces a homomorphism

$$K(\text{Var})/(\mathbb{L}) \rightarrow \mathbb{Z}[u, v]/(uv).$$

The image of $\mathcal{R}/(\mathbb{L}) \cong \mathbb{Z}$ in $\mathbb{Z}[u, v]/(uv)$ is \mathbb{Z} , and the statement follows. □

For example, $h^{1,0}(X) \neq 0$ for e.g., a smooth elliptic curve. Therefore, Corollary 3.8 shows that classes of elliptic curves are not in the span of the classes $[Y_G]$.

References

1. Abramovich, D., Karu, K., Matsuki, K., Włodarczyk, J.: Torification and factorization of birational maps. *J. Am. Math. Soc.* **15**, 531–572 (2002)
2. Aluffi, P., Marcolli, M.: Feynman motives of banana graphs. *Commun. Number Theory Phys.* **3**, 1–57 (2009)
3. Aluffi, P., Marcolli, M.: Algebraic-geometric Feynman rules. *Int. J. Geom. Methods Mod. Phys.* (to appear). arXiv:0811.2514
4. Aluffi, P., Marcolli, M.: Feynman motives and deletion–contraction relations. *Topology of Algebraic Varieties and Singularities, Proceedings of the Lib60ber 2009 Jaca Conference* (to appear). arXiv:0907.3225
5. Bittner, F.: The universal Euler characteristic for varieties of characteristic zero. *Compos. Math.* **140**(4), 1011–1032 (2004)
6. Belkale, P., Brosnan, P.: Matroids, motives, and a conjecture of Kontsevich. *Duke Math. J.* **116**, 147–188 (2003)
7. Bloch, S., Esnault, H., Kreimer, D.: On motives associated to graph polynomials. *Commun. Math. Phys.* **267**(1), 181–225 (2006)
8. Brown, F., Schnetz, O.: A K3 in ϕ^4 . arXiv:1006.4064
9. Danilov, V.I., Khovanskii, A.G.: Newton polyhedra and an algorithm for calculating Hodge–Deligne numbers. *Izv. Akad. Nauk SSSR Ser. Mat.* **50**(5), 925–945 (1986)

10. Greco, S., Vistoli, A.: On the construction of rational surfaces with assigned singularities. In: *Algebraic Geometry—Open Problems (Ravello, 1982)*. Lecture Notes in Mathematics, vol. 997, pp. 210–217. Springer, Berlin (1983)
11. Kollár, J.: Conics in the Grothendieck ring. *Adv. Math.* **198**(1), 27–35 (2005)
12. Larsen, M., Lunts, V.A.: Motivic measures and stable birational geometry. *Mosc. Math. J.* **3**(1), 85–95, 259 (2003)
13. Schnetz, O.: Quantum field theory over \mathbb{F}_q . arXiv:0909.0905
14. Stembridge, J.R.: Counting points on varieties over finite fields related to a conjecture of Kontsevich. *Ann. Comb.* **2**(4), 365–385 (1998)
15. Veys, W.: Arc spaces, motivic integration and stringy invariants. In: “Singularity Theory and its Applications, vol. 43, pp. 529–572. *Adv. Stud. Pure Math. Math. Soc.*, Japan (2006)