

# Some remarks concerning Voevodsky’s nilpotence conjecture

By *Marcello Bernardara* at Toulouse, *Matilde Marcolli* at Pasadena and  
*Gonçalo Tabuada* at Cambridge, MA

---

**Abstract.** In this article we extend Voevodsky’s nilpotence conjecture from smooth projective schemes to the broader setting of smooth proper dg categories. Making use of this noncommutative generalization, we then address Voevodsky’s original conjecture in the following cases: quadric fibrations, intersection of quadrics, linear sections of Grassmannians, linear sections of determinantal varieties, homological projective duals, and Moishetzon manifolds.

## 1. Introduction and statement of results

Let  $k$  be a base field and  $F$  a field of coefficients of characteristic zero.

**Voevodsky’s nilpotence conjecture.** In a foundational work [36], Voevodsky introduced the smash-nilpotence equivalence relation  $\sim_{\otimes\text{nil}}$  on algebraic cycles and conjectured its agreement with the classical numerical equivalence relation  $\sim_{\text{num}}$ . Concretely, given a smooth projective  $k$ -scheme  $X$ , he stated the following:

**Conjecture  $V(X)$ .** We have  $\mathcal{Z}^*(X)_F / \sim_{\otimes\text{nil}} = \mathcal{Z}^*(X)_F / \sim_{\text{num}}$ .

Thanks to the work of Kahn–Sebastian, Matsusaka, Voevodsky, and Voisin (see, for instance, [16, 30, 36, 37] and also [1, Section 11.5.2.3]), the above conjecture holds in the case of curves, surfaces, and abelian 3-folds (when  $k$  is of characteristic zero).

---

G. Tabuada was partially supported by the National Science Foundation CAREER Award #1350472 and by the Fundação para a Ciência e a Tecnologia (Portuguese Foundation for Science and Technology) through the project grant UID/MAT/00297/2013 (Centro de Matemática e Aplicações). M. Marcolli was partially supported by the NSF grants DMS-1201512 and PHY-1205440.

Sections 3–5 are a revised version of half of an old preprint [26] by M. Marcolli and G. Tabuada. The remaining work is a recent collaboration between M. Bernardara and G. Tabuada.

**Noncommutative nilpotence conjecture.** A dg category  $\mathcal{A}$  is a category enriched over dg  $k$ -vector spaces; see Section 2.1. Following Kontsevich [18–20],  $\mathcal{A}$  is called *smooth* if it is perfect as a bimodule over itself and *proper* if for any two objects  $x, y \in \mathcal{A}$  we have  $\sum_i \dim H^i \mathcal{A}(x, y) < \infty$ . The classical example is the unique dg enhancement  $\text{perf}_{\text{dg}}(X)$  of the category of perfect complexes  $\text{perf}(X)$  of a smooth projective  $k$ -scheme  $X$ ; see Lunts and Orlov [25]. As explained in Section 2.3–2.4, the Grothendieck group  $K_0(\mathcal{A})$  of every smooth proper dg category  $\mathcal{A}$  comes endowed with a  $\otimes$ -nilpotence equivalence relation  $\sim_{\otimes\text{nil}}$  and with a numerical equivalence relation  $\sim_{\text{num}}$ . Motivated by the above conjecture, we state the following:

**Conjecture  $V_{\text{NC}}(\mathcal{A})$ .** We have  $K_0(\mathcal{A})_F / \sim_{\otimes\text{nil}} = K_0(\mathcal{A})_F / \sim_{\text{num}}$ .

Our first main result is the following reformulation of Voevodsky’s conjecture:

**Theorem 1.1.** *Conjecture  $V(X)$  is equivalent to Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(X))$ .*

Theorem 1.1 shows us that when restricted to the commutative world, the noncommutative nilpotence conjecture reduces to Voevodsky’s original conjecture. Making use of this noncommutative viewpoint, we now address Voevodsky’s nilpotence conjecture in several cases.

**Quadric fibrations.** Let  $S$  be a smooth projective  $k$ -scheme and  $q : Q \rightarrow S$  a flat quadric fibration of relative dimension  $n$  with  $Q$  smooth. Recall from Kuznetsov [23] (see also [5]) the construction of the sheaf  $\mathcal{C}_0$  of even parts of the Clifford algebra associated to  $q$ . Recall also from [23] that when the discriminant divisor of  $q$  is smooth and  $n$  is even (resp. odd), we have a discriminant double cover  $\tilde{S} \rightarrow S$  (resp. a square root stack  $\widehat{S}$ ) equipped with an Azumaya algebra  $\mathcal{B}_0$ . Our second main result allows us to decompose Conjecture  $V(Q)$  into simpler pieces:

**Theorem 1.2.** *The following hold:*

(i) *We have*

$$V(Q) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(S, \mathcal{C}_0)) + V(S).$$

(ii) *When the discriminant divisor of  $q$  is smooth and  $n$  is even, we have*

$$V(Q) \iff V(\tilde{S}) + V(S).$$

*As a consequence,  $V(Q)$  holds when  $\dim(S) \leq 2$ , and becomes equivalent to  $V(\tilde{S})$  when  $S$  is an abelian 3-fold and  $k$  is of characteristic zero.*

(iii) *When the discriminant divisor of  $q$  is smooth and  $n$  is odd, we have*

$$V(Q) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{S}, \mathcal{B}_0)) + V(S).$$

*As a consequence, Conjecture  $V(Q)$  becomes equivalent to  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{S}), \mathcal{B}_0)$  when  $\dim(S) \leq 2$ . This latter conjecture holds when  $\dim(S) \leq 2$ .*

**Remark 1.3.** The (rational) Chow motive of a quadric fibration  $q : Q \rightarrow S$  was computed by Vial in [35, Theorem 4.2 and Corollary 4.4]. In the particular case where  $\dim(S) \leq 2$ ,

it consists of a direct sum of submotives of smooth projective  $k$ -schemes of dimension at most two. This motivic decomposition provides an alternative “geometric” proof of Conjecture  $V(Q)$ . We will rely on this argument to prove the last statement of item (iii). In the particular case where  $S$  is a curve and  $k$  is algebraically closed, we provide also a “categorical” proof of this last statement; see Remark 6.5. The fact that  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{S}), \mathcal{B}_0)$  holds when  $\dim(S) \leq 2$  will play a key role in the proof of Theorem 1.4 below.

**Intersection of quadrics.** Let  $X$  be a smooth complete intersection of  $r$  quadric hypersurfaces in  $\mathbb{P}^m$ . The linear span of these  $r$  quadrics gives rise to a hypersurface  $Q \subset \mathbb{P}^{r-1} \times \mathbb{P}^m$ , and the projection into the first factor is a flat quadric fibration  $q : Q \rightarrow \mathbb{P}^{r-1}$  of relative dimension  $m - 1$ .

**Theorem 1.4.** *The following holds:*

(i) *We have*

$$V(X) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0)).$$

(ii) *When the discriminant divisor of  $q$  is smooth and  $m$  is odd, we have*

$$V(X) \iff V(\widehat{\mathbb{P}^{r-1}}).$$

*As a consequence,  $V(X)$  holds when  $r \leq 3$ .*

(iii) *When the discriminant divisor of  $q$  is smooth and  $m$  is even, we have*

$$V(X) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{\mathbb{P}^{r-1}}, \mathcal{B}_0)).$$

*This latter conjecture holds when  $r \leq 3$  and  $k$  is algebraically closed.*

**Remark 1.5.** The (rational) Chow motive of a complete intersection  $X$  was computed in [8, Theorem 2.1] in the particular cases where  $r \leq 2$  or  $r = 3$  and  $m$  is odd. It consists of a direct sum of submotives of smooth projective  $k$ -schemes of dimension at most one. This motivic decomposition provides an alternative proof of Conjecture  $V(X)$ . A similar argument holds in the case where  $r = 3$  and  $m$  is even.

**Remark 1.6** (Relative version). Theorem 1.4 has a relative analogue with  $X$  replaced by a generic relative complete intersection  $X \rightarrow S$  of  $r$  quadric fibrations  $Q_i \rightarrow S$  of relative dimension  $m - 1$ ; consult [5, Definition 1.2.4] for details. Items (i), (ii), and (iii), hold similarly with  $\mathbb{P}^{r-1}$  replaced by a  $\mathbb{P}^{r-1}$ -bundle  $T \rightarrow S$ , with  $V(\widehat{\mathbb{P}^{r-1}})$  replaced by  $V(\widehat{T}) + V(S)$ , and with  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{\mathbb{P}^{r-1}}, \mathcal{B}_0))$  replaced by  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\widehat{T}, \mathcal{B}_0)) + V(S)$ , respectively. Note that thanks to the relative item (ii), Conjecture  $V(X)$  holds when  $r = 2$  and  $S$  is a curve.

**Linear sections of Grassmannians.** Following Kuznetsov [21], consider the following two classes of schemes:

(i) Let  $X_L$  be a generic linear section of codimension  $r$  of the Grassmannian  $\text{Gr}(2, W)$  (with  $W = k^{\oplus 6}$ ) under the Plücker embedding, and  $Y_L$  the corresponding dual linear section of the cubic Pfaffian  $\text{Pf}(4, W^*)$  in  $\mathbb{P}(\Lambda^2 W^*)$ .

For example when  $r = 3$ ,  $X_L$  is a Fano 5-fold; when  $r = 4$ ,  $X_L$  is a Fano 4-fold; and when  $r = 6$ ,  $X_L$  is a  $K3$  surface of degree 14 and  $Y_L$  a Pfaffian cubic 4-fold. Moreover,  $X_L$  and  $Y_L$  are smooth whenever  $r \leq 6$ .

- (ii) Let  $X_L$  be a generic linear section of codimension  $r$  of the Grassmannian  $\text{Gr}(2, W)$  (with  $W = k^{\oplus 7}$ ) under the Plücker embedding, and  $Y_L$  the corresponding dual linear section of the cubic Pfaffian  $\text{Pf}(4, W^*)$  in  $\mathbb{P}(\Lambda^2 W^*)$ .

For example when  $r = 5$ ,  $X_L$  is a Fano 5-fold; when  $r = 4$ ,  $X_L$  is a Fano 4-fold; when  $r = 8$ ,  $Y_L$  is a Fano 4-fold; and when  $r = 9$ ,  $Y_L$  is a Fano 5-fold. Moreover,  $X_L$  and  $Y_L$  are smooth whenever  $r \leq 10$ .

**Theorem 1.7.** *Let  $X_L$  and  $Y_L$  be as in the above classes (i)–(ii). Under the assumption that  $X_L$  and  $Y_L$  are smooth, we have*

$$V(X_L) \iff V(Y_L).$$

*This conjecture holds when  $r \leq 6$  (class (i)), and when  $r \leq 6$  and  $8 \leq r \leq 10$  (class (ii)).*

**Remark 1.8.** To the best of the authors’ knowledge, Theorem 1.7 proves Voevodsky’s nilpotence conjecture in new cases.

**Linear sections of determinantal varieties.** Let  $U$  and  $V$  be  $k$ -vector spaces of dimensions  $m$  and  $n$ , respectively, with  $n \geq m$ , and  $0 < r < m$  an integer. Following [7], consider the universal determinantal variety  $Z_{m,n}^r \subset \mathbb{P}(U \otimes V)$  given by the locus of matrices  $M : U^* \rightarrow V$  of rank at most  $r$ . Its Springer resolution is denoted by

$$\mathcal{X}_{m,n}^r := \mathbb{P}(\mathcal{U} \otimes V) \rightarrow \text{Gr}(r, U),$$

where  $\mathcal{U}$  stands for the tautological subbundle on  $\text{Gr}(r, U)$ . Under these notations, we have the following class of schemes:

- (i) Let  $X_L$  be a generic linear section of codimension  $c$  of  $\mathcal{X}_{m,n}^r$  under the map

$$\mathcal{X}_{m,n}^r \rightarrow \mathbb{P}(U \otimes V),$$

and  $Y_L$  the corresponding dual linear section of  $\mathcal{X}_{m,n}^{m-r}$  under the map

$$\mathcal{X}_{m,n}^{m-r} \rightarrow \mathbb{P}(U^* \otimes V^*).$$

**Remark 1.9.** As explained in [7, Section 3],  $X_L$  and  $Y_L$  are smooth crepant categorical resolution of singularities of  $(Z_{m,n}^r)_L$  and  $(Z_{m,n}^{m-r})_L$ , respectively.

For example when  $m = n = 4$  and  $r = 1$ ,  $X_L$  is a  $(6 - c)$ -dimensional section of  $\mathbb{P}^3 \times \mathbb{P}^3$  under the Segre embedding, and  $Y_L$  the resolution of the dual  $(c - 2)$ -dimensional determinantal quartic. In the same vein, when  $m = n = 4$  and  $r = 2$ ,  $X_L$  is an  $(11 - c)$ -dimensional section of the self-dual orbit of  $4 \times 4$  matrices of rank 2, and  $Y_L$  the  $(c - 5)$ -dimensional dual section. Moreover, the following holds:

- (a) When  $c \leq 7$ ,  $X_L$  is a Fano  $(11 - c)$ -fold and  $\dim(Y_L) \leq 2$ .
- (b) When  $c = 8$ ,  $X_L$  and  $Y_L$  are dual Calabi–Yau 3-folds.
- (c) When  $c \geq 9$ ,  $Y_L$  is a Fano  $(c - 5)$ -fold and  $\dim(X_L) \leq 2$ .

For further example, consult [7, Section 3.3] and well as [7, Tables 1 and 2].

**Theorem 1.10.** *When  $X_L$  and  $Y_L$  are as in the above class (i), we have*

$$V(X_L) \iff V(Y_L).$$

*This conjecture holds when  $\dim(X_L) \leq 2$  or  $\dim(Y_L) \leq 2$ .*

**Remark 1.11.** To the best of the authors’ knowledge, Theorem 1.10 proves Voevodsky’s nilpotence conjecture in new cases.

**Homological projective duality.** Making use of Kuznetsov’s theory of homological projective duality (HPD) [22], Theorem 1.7 admits the following generalization: let  $X$  be a smooth projective  $k$ -scheme equipped with an ample line bundle  $\mathcal{O}_X(1)$ . Note that  $\mathcal{O}_X(1)$  gives rise to a morphism  $X \rightarrow \mathbb{P}(V)$ , where  $V := H^0(X, \mathcal{O}_X(1))^*$ . Let  $Y$  be the HP-dual of  $X$ ,  $\mathcal{O}_Y(1)$  the associated ample line bundle, and  $Y \rightarrow \mathbb{P}(V^*)$  the associated morphism. Assume that  $\text{perf}(X)$  admits a *Lefschetz decomposition*  $\text{perf}(X) = \langle \mathbb{A}_0, \dots, \mathbb{A}_n(n) \rangle$  with respect to  $\mathcal{O}_X(1)$ , i.e. a semi-orthogonal decomposition of  $\text{perf}(X)$  such that  $\mathbb{A}_0 \supset \dots \supset \mathbb{A}_n$  and  $\mathbb{A}_i(i) := \mathbb{A}_i \otimes \mathcal{O}(i)$ . Assume also that Conjecture  $V_{\text{NC}}(\mathbb{A}_0^{\text{dg}})$  holds, where  $\mathbb{A}_0^{\text{dg}}$  stands for the dg enhancement of  $\mathbb{A}_0$  induced from  $\text{perf}_{\text{dg}}(X)$ ; see Section 2.2. Finally, let  $L \subset V$  be a subspace such that the linear sections  $X_L := X \times_{\mathbb{P}(V)} \mathbb{P}(L)$  and  $Y_L := Y \times_{\mathbb{P}(V^*)} \mathbb{P}(L^\perp)$  are of expected dimension  $\dim(X_L) = \dim(X) - \dim(L)$  and  $\dim(Y_L) = \dim(Y) - \dim(L^\perp)$ , respectively.

**Theorem 1.12.** *Let  $X_L$  and  $Y_L$  be as above. Under the assumption that  $X_L$  and  $Y_L$  are smooth, we have*

$$V(X_L) \iff V(Y_L).$$

**Remark 1.13.** Theorem 1.12 reduces to Theorem 1.7 (resp. to Theorem 1.10) in the particular case of Grassmannian–Pfaffian (resp. determinantal) homological projective duality; consult [21] (resp. [7]) for details.

**Moishezon manifolds.** A *Moishezon manifold*  $X$  is a compact complex manifold such that the field of meromorphic functions on each component of  $X$  has transcendence degree equal to the dimension of the component. As proved by Moishezon [31],  $X$  is a smooth projective  $\mathbb{C}$ -scheme if and only if it admits a Kähler metric. In the remaining cases, Artin [3] showed that  $X$  is a proper algebraic space over  $\mathbb{C}$ .

Let  $Y \rightarrow \mathbb{P}^2$  be one of the non-rational conic bundles described by Artin and Mumford in [4], and  $X \rightarrow Y$  a small resolution. In this case,  $X$  is a smooth (not necessarily projective) Moishezon manifold.

**Theorem 1.14.** *Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(X))$  holds for the above resolutions.*

**Remark 1.15.** The proofs of Theorems 1.2, 1.4, 1.7, 1.10, 1.12, and 1.14 are based on the study of a smooth projective  $k$ -scheme (or algebraic space)  $X$  via semi-orthogonal decompositions of its category of perfect complexes  $\text{perf}(X)$ ; see Bondal–Orlov [9], Huybrechts [14], and Kuznetsov [24] for instance. This approach allows the reduction of Voevodsky’s Conjecture  $V(X)$  to several noncommutative conjectures  $V_{\text{NC}}$  – one for each piece of the semi-orthogonal decomposition. We believe this provides a new tool for the proof of Voevodsky’s conjecture as well as of its generalization to algebraic spaces.

## 2. Preliminaries

**2.1. Dg categories.** A differential graded category (in short, dg category)  $\mathcal{A}$  is a category enriched over dg  $k$ -vector spaces; consult Keller [17] for details. For example, every (dg)  $k$ -algebra  $A$  gives naturally rise to a dg category  $\underline{A}$  with a single object. Let  $\text{dgcats}$  be the category of small dg categories. Recall from [17, Section 3] the construction of the derived category  $\mathcal{D}(\mathcal{A})$  of  $\mathcal{A}$ . This triangulated category admits arbitrary direct sums and we will write  $\mathcal{D}_c(\mathcal{A})$  for the full subcategory of compact objects. A dg functor  $\mathcal{A} \rightarrow \mathcal{B}$  is called a *Morita equivalence* if it induces an equivalence  $\mathcal{D}(\mathcal{A}) \xrightarrow{\sim} \mathcal{D}(\mathcal{B})$ . Finally, let us write  $\mathcal{A} \otimes \mathcal{B}$  for the tensor product of dg categories.

**2.2. Perfect complexes.** Given a stack  $\mathcal{X}$  and a sheaf of  $\mathcal{O}_{\mathcal{X}}$ -algebras  $\mathcal{G}$ , let  $\text{Mod}(\mathcal{X}, \mathcal{G})$  be the Grothendieck category of sheaves of (right)  $\mathcal{G}$ -modules,  $\mathcal{D}(\mathcal{X}, \mathcal{G}) := \mathcal{D}(\text{Mod}(\mathcal{X}, \mathcal{G}))$  the derived category of  $\mathcal{G}$ , and  $\text{perf}(\mathcal{X}, \mathcal{G})$  the subcategory of perfect complexes. As explained in [17, Section 4.4], the derived category  $\mathcal{D}_{\text{dg}}(\mathcal{E}x)$  of an abelian (or exact) category  $\mathcal{E}x$  is defined as the (Drinfeld’s) dg quotient  $\mathcal{C}_{\text{dg}}(\mathcal{E}x)/\text{Ac}_{\text{dg}}(\mathcal{E}x)$  of the dg category of complexes over  $\mathcal{E}x$  by its full dg subcategory of acyclic complexes. Hence, let us write  $\mathcal{D}_{\text{dg}}(\mathcal{X}, \mathcal{G})$  for the dg category  $\mathcal{D}_{\text{dg}}(\mathcal{E}x)$  with  $\mathcal{E}x := \text{Mod}(\mathcal{X}, \mathcal{G})$  and  $\text{perf}_{\text{dg}}(\mathcal{X}, \mathcal{G})$  for the full dg subcategory of perfect complexes.

**Lemma 2.1.** *Let  $X$  be a smooth projective  $k$ -scheme, and let  $\text{perf}(X) = \langle \mathcal{T}_1, \dots, \mathcal{T}_n \rangle$  be a semi-orthogonal decomposition. In this case, the dg categories  $\mathcal{T}_i^{\text{dg}}$  (where  $\mathcal{T}_i^{\text{dg}}$  stands for the dg enhancement of  $\mathcal{T}_i$  induced from  $\text{perf}_{\text{dg}}(X)$ ) are smooth and proper.*

*Proof.* Let  $\text{Ho}(\text{dgcats})$  be the localization of  $\text{dgcats}$  with respect to the class of Morita equivalences. The tensor product of dg categories gives rise to a symmetric monoidal structure on  $\text{dgcats}$  which descends to  $\text{Ho}(\text{dgcats})$ . Moreover, as proved in [10, Theorem 5.8], the smooth and proper dg categories can be characterized as those objects of  $\text{Ho}(\text{dgcats})$  which are dualizable. Note that the canonical inclusion  $\mathcal{T}_i^{\text{dg}} \hookrightarrow \text{perf}_{\text{dg}}(X)$  and projection  $\text{perf}_{\text{dg}}(X) \rightarrow \mathcal{T}_i^{\text{dg}}$  dg functors express  $\mathcal{T}_i^{\text{dg}}$  as a direct factor of  $\text{perf}_{\text{dg}}(X)$  in  $\text{Ho}(\text{dgcats})$ . Hence, since  $\text{perf}_{\text{dg}}(X)$  is smooth and proper, we conclude that  $\mathcal{T}_i^{\text{dg}}$  is also smooth and proper.  $\square$

**2.3.  $\otimes$ -nilpotence equivalence relation.** Let  $\mathcal{A}$  be a dg category. An element  $[M]$  of the Grothendieck group  $K_0(\mathcal{A}) := K_0(\mathcal{D}_c(\mathcal{A}))$  is called  $\otimes$ -nilpotent if there exists an integer  $n > 0$  such that  $[M^{\otimes n}] = 0$  in  $K_0(\mathcal{A}^{\otimes n})$ . This gives rise to a well-defined equivalence relation  $\sim_{\otimes\text{nil}}$  on  $K_0(\mathcal{A})$  and on its  $F$ -linearization  $K_0(\mathcal{A})_F$ .

**2.4. Numerical equivalence relation.** Let  $\mathcal{A}$  be a smooth proper dg category. As explained in [27, Section 4], the pairing  $(M, N) \mapsto \sum_i (-1)^i \dim \text{Hom}_{\mathcal{D}_c(\mathcal{A})}(M, N[i])$  gives rise to a well-defined bilinear form  $\chi(-, -)$  on  $K_0(\mathcal{A})$ . Moreover, the left and right kernels of  $\chi(-, -)$  are the same. An element  $[M]$  of the Grothendieck group  $K_0(\mathcal{A})$  is said to be *numerically trivial* if  $\chi([M], [N]) = 0$  for all elements  $[N] \in K_0(\mathcal{A})$ . This gives rise to an equivalence relation  $\sim_{\text{num}}$  on  $K_0(\mathcal{A})$  and consequently on  $K_0(\mathcal{A})_F$ . When  $\mathcal{A} = \text{perf}_{\text{dg}}(X)$ , with  $X$  a smooth projective  $k$ -scheme, and  $F = \mathbb{Q}$ , this equivalence relation reduces, via the Chern character  $K_0(X)_{\mathbb{Q}} \xrightarrow{\sim} CH^*(X)_{\mathbb{Q}}$ , to the classical numerical equivalence relation on the Chow ring  $CH^*(X)_{\mathbb{Q}}$ .

**2.5. Motives.** We assume the reader is familiar with the categories of Chow motives  $\text{Chow}(k)_F$  and numerical motives  $\text{Num}(k)_F$ ; see [1, Chapitre 4]. The Tate motive will be denoted  $F(1)$ . In the same vein, we assume some familiarity with the categories of noncommutative Chow motives  $\text{NChow}(k)_F$  and noncommutative numerical motives  $\text{NNum}(k)_F$ ; consult the surveys [28, Sections 2–3] and [32, Section 4], and the references therein. Recall from [1, Chapitre 4] that  $\text{NNum}(k)_F$  is the idempotent completion of the quotient of  $\text{NChow}(k)_F$  by its largest  $\otimes$ -ideal<sup>1)</sup>, and that  $\text{Hom}_{\text{NChow}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F$ .

### 3. Orbit categories and $\otimes$ -nilpotence

Let  $\mathcal{C}$  be an  $F$ -linear additive rigid symmetric monoidal category.

**Orbit categories.** Given a  $\otimes$ -invertible object  $\mathcal{O} \in \mathcal{C}$ , recall from [33, Section 7] the construction of the orbit category  $\mathcal{C}/_{\otimes \mathcal{O}}$ . It has the same objects as  $\mathcal{C}$  and morphisms

$$\text{Hom}_{\mathcal{C}/_{\otimes \mathcal{O}}}(a, b) := \bigoplus_{j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(a, b \otimes \mathcal{O}^{\otimes j}).$$

The composition law is induced from  $\mathcal{C}$ . By construction,  $\mathcal{C}/_{\otimes \mathcal{O}}$  is  $F$ -linear, additive, and comes equipped with a canonical projection functor  $\pi : \mathcal{C} \rightarrow \mathcal{C}/_{\otimes \mathcal{O}}$ . Moreover,  $\pi$  is endowed with a natural 2-isomorphism  $\pi \circ (- \otimes \mathcal{O}) \xrightarrow{\sim} \pi$  and is 2-universal among all such functors. As proved in [33, Lemma 7.3],  $\mathcal{C}/_{\otimes \mathcal{O}}$  inherits from  $\mathcal{C}$  a symmetric monoidal structure making  $\pi$  symmetric monoidal. On objects it is the same. On morphisms it is defined as the unique bilinear pairing

$$\bigoplus_{j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(a, b \otimes \mathcal{O}^{\otimes j}) \times \bigoplus_{j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(c, d \otimes \mathcal{O}^{\otimes j}) \rightarrow \bigoplus_{j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(a \otimes c, (b \otimes d) \otimes \mathcal{O}^{\otimes j})$$

which sends the pair

$$(a \xrightarrow{f_r} b \otimes \mathcal{O}^{\otimes r}, c \xrightarrow{g_s} d \otimes \mathcal{O}^{\otimes s})$$

to

$$(f \otimes g)_{(r+s)} : a \otimes c \xrightarrow{f_r \otimes g_s} b \otimes \mathcal{O}^{\otimes r} \otimes d \otimes \mathcal{O}^{\otimes s} \simeq (b \otimes d) \otimes \mathcal{O}^{\otimes (r+s)}.$$

**$\otimes$ -nilpotence.** The  $\otimes_{\text{nil}}$ -ideal of  $\mathcal{C}$  is defined as

$$\otimes_{\text{nil}}(a, b) := \{f \in \text{Hom}_{\mathcal{C}}(a, b) : f^{\otimes n} = 0 \text{ for } n \gg 0\}.$$

By construction,  $\otimes_{\text{nil}}$  is a  $\otimes$ -ideal. Moreover, all its ideals  $\otimes_{\text{nil}}(a, a) \subset \text{Hom}_{\mathcal{C}}(a, a)$  are nilpotent; see [2, Lemma 7.4.2 (ii)]. As a consequence, the  $\otimes$ -functor  $\mathcal{C} \rightarrow \mathcal{C}/_{\otimes_{\text{nil}}}$  is not only  $F$ -linear and additive but moreover conservative. Furthermore, since idempotents can be lifted along nilpotent ideals (see [6, Section III, Proposition 2.10]),  $\mathcal{C}/_{\otimes_{\text{nil}}}$  is idempotent complete whenever  $\mathcal{C}$  is idempotent complete.

<sup>1)</sup> Different from the entire category  $\text{NChow}(k)_F$ .

**Compatibility.** Let  $\mathcal{C}$  be a category and  $\mathcal{O} \in \mathcal{C}$  a  $\otimes$ -invertible objects as above.

**Proposition 3.1.** *There exists a canonical  $F$ -linear additive  $\otimes$ -equivalence  $\theta$  making the following diagram commute:*

$$(3.2) \quad \begin{array}{ccc} \mathcal{C}/\otimes_{\text{nil}} & \longleftarrow \mathcal{C} & \longrightarrow \mathcal{C}/\otimes \mathcal{O} \\ \downarrow & & \downarrow \\ (\mathcal{C}/\otimes_{\text{nil}})/\otimes \mathcal{O} & \xrightarrow[\theta]{\cong} & (\mathcal{C}/\otimes \mathcal{O})/\otimes_{\text{nil}}. \end{array}$$

*Proof.* The existence of the  $F$ -linear additive  $\otimes$ -functor  $\theta$  follows from the fact that

$$(3.3) \quad \mathcal{C} \longrightarrow \mathcal{C}/\otimes \mathcal{O} \longrightarrow (\mathcal{C}/\otimes \mathcal{O})/\otimes_{\text{nil}}$$

vanishes on the  $\otimes_{\text{nil}}$ -ideal and also from the natural 2-isomorphism between (3.3)  $\circ$   $(-\otimes \mathcal{O})$  and (3.3). Note that the functor  $\theta$  is the identity on objects and sends  $\{[f_j]\}_{j \in \mathbb{Z}}$  to  $\{[f_j]\}_{j \in \mathbb{Z}}$ . Clearly, it is full. The faithfulness is left as an exercise.  $\square$

#### 4. $\otimes$ -nilpotence of motives

By construction, the categories  $\text{Chow}(k)_F$  and  $\text{NChow}(k)_F$  are  $F$ -linear, additive, rigid symmetric monoidal, and idempotent complete. Let us denote by

$$\text{Voev}(k)_F := \text{Chow}(k)_F / \otimes_{\text{nil}}$$

and

$$\text{NVoev}(k)_F := \text{NChow}(k)_F / \otimes_{\text{nil}}$$

the associated quotients. They fit in the following sequences:

$$\text{Chow}(k)_F \rightarrow \text{Voev}(k)_F \rightarrow \text{Num}(k)_F$$

and

$$\text{NChow}(k)_F \rightarrow \text{NVoev}(k)_F \rightarrow \text{NNum}(k)_F.$$

The relation between all these motivic categories is the following:

**Proposition 4.1.** *There exist  $F$ -linear additive fully-faithful  $\otimes$ -functors  $R, R_{\otimes_{\text{nil}}}, R_{\mathcal{N}}$  making the following diagram commute:*

$$(4.2) \quad \begin{array}{ccccc} \text{Chow}(k)_F & \xrightarrow{\pi} & \text{Chow}(k)_F / \otimes_{F(1)} & \xrightarrow{R} & \text{NChow}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Voev}(k)_F & \xrightarrow{\pi} & \text{Voev}(k)_F / \otimes_{F(1)} & \xrightarrow{R_{\otimes_{\text{nil}}}} & \text{NVoev}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Num}(k)_F & \xrightarrow{\pi} & \text{Num}(k)_F / \otimes_{F(1)} & \xrightarrow{R_{\mathcal{N}}} & \text{NNum}(k)_F. \end{array}$$

*Proof.* The outer commutative square, with  $R, R_{\mathcal{N}}$  both  $F$ -linear additive fully-faithful  $\otimes$ -functors, was built in [29, Theorem 1.13]. Consider now the “zoomed” diagram

$$(4.3) \quad \begin{array}{ccccc} \text{Chow}(k)_F / \otimes_{F(1)} & \xlongequal{\quad} & \text{Chow}(k)_F / \otimes_{F(1)} & \xrightarrow{R} & \text{NChow}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Voev}(k)_F / \otimes_{F(1)} & \xrightarrow[\theta]{\cong} & (\text{Chow}(k)_F / \otimes_{F(1)}) / \otimes_{\text{nil}} & \xrightarrow{R / \otimes_{\text{nil}}} & \text{NVoev}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Num}(k)_F / \otimes_{F(1)} & \xlongequal{\quad} & \text{Num}(k)_F / \otimes_{F(1)} & \xrightarrow{R_{\mathcal{N}}} & \text{NNum}(k)_F \end{array}$$

By definition,  $R_{\otimes_{\text{nil}}} := R / \otimes_{\text{nil}} \circ \theta$ . Since  $R$  is an  $F$ -linear additive fully-faithful  $\otimes$ -functor, we conclude that  $R_{\otimes_{\text{nil}}}$  is also an  $F$ -linear additive fully-faithful  $\otimes$ -functor. The commutativity of the bottom squares of diagram (4.3) follows from the fact that  $\text{Num}(k)_F / \otimes_{F(1)}$  identifies with the quotient of  $\text{Chow}(k)_F / \otimes_{F(1)}$  by its largest  $\otimes$ -ideal  $\mathcal{N}$ ; consult [29, Proposition 3.2] for details.  $\square$

### 5. Proof of Theorem 1.1

Note first that we have the following natural isomorphisms:

$$\begin{aligned} \text{Hom}_{\text{Voev}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X) &\simeq \mathcal{Z}^*(X)_F / \sim_{\otimes_{\text{nil}}}, \\ \text{Hom}_{\text{Num}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X) &\simeq \mathcal{Z}^*(X)_F / \sim_{\text{num}}. \end{aligned}$$

As a consequence, Conjecture  $V(X)$  becomes equivalent to the injectivity of

$$(5.1) \quad \text{Hom}_{\text{Voev}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X) \twoheadrightarrow \text{Hom}_{\text{Num}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X).$$

Given a smooth and proper dg category  $\mathcal{A}$ , we have also natural isomorphisms

$$\text{Hom}_{\text{NVoev}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F / \sim_{\otimes_{\text{nil}}}, \quad \text{Hom}_{\text{NNum}(k)_F}(\underline{k}, \mathcal{A}) \simeq K_0(\mathcal{A})_F / \sim_{\text{num}}.$$

Hence, Conjecture  $V_{\text{NC}}(\mathcal{A})$  becomes equivalent to the injectivity of

$$(5.2) \quad \text{Hom}_{\text{NVoev}(k)_F}(\underline{k}, \mathcal{A}) \twoheadrightarrow \text{Hom}_{\text{NNum}(k)_F}(\underline{k}, \mathcal{A}).$$

Now, recall from [33, Theorem 1.1] that the image of  $X$  under the composed functor  $R \circ \pi$  identifies naturally with the noncommutative Chow motive  $\text{perf}_{\text{dg}}(X)$ . Similarly, the image of  $\text{Spec}(k)$  under  $R \circ \pi$  identifies with  $\text{perf}_{\text{dg}}(\text{Spec}(k))$  which is Morita equivalent to  $\underline{k}$ . As a consequence, since the functors  $R_{\otimes_{\text{nil}}}$  and  $R_{\mathcal{N}}$  are fully-faithful, the bottom right-hand side square of diagram (4.2) gives rise to the following commutative diagram:

$$\begin{array}{ccc} \text{Hom}_{\text{Voev}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X) & \xrightarrow{\cong} & \text{Hom}_{\text{NVoev}(k)_F}(\underline{k}, \text{perf}_{\text{dg}}(X)) \\ \downarrow (5.1) & & \downarrow (5.2) \\ \text{Hom}_{\text{Num}(k)_F / \otimes_{F(1)}}(\text{Spec}(k), X) & \xrightarrow{\cong} & \text{Hom}_{\text{NNum}(k)_F}(\underline{k}, \text{perf}_{\text{dg}}(X)). \end{array}$$

Using the above reformulations of Conjectures  $V$  and  $V_{\text{NC}}$ , we conclude finally that Conjecture  $V(X)$  is equivalent to Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(X))$ .

### 6. Proof of Theorem 1.2

**Item (i).** As proved by Kuznetsov in [23, Theorem 4.2], one has the semi-orthogonal decomposition

$$\text{perf}(Q) = \langle \text{perf}(S, \mathcal{C}_0), \text{perf}(S)_1, \dots, \text{perf}(S)_n \rangle$$

with  $\text{perf}(S)_i := q^* \text{perf}(S) \otimes \mathcal{O}_{Q/S}(i)$ . Note that we have  $\text{perf}(S)_i \simeq \text{perf}(S)$  for every  $i$ . Using [8, Proposition 3.1], one then obtains a direct sum decomposition in  $\text{NChow}(k)_F$

$$(6.1) \quad \text{perf}_{\text{dg}}(Q) \simeq \text{perf}^{\text{dg}}(S, \mathcal{C}_0) \oplus \underbrace{\text{perf}_{\text{dg}}(S) \oplus \dots \oplus \text{perf}_{\text{dg}}(S)}_{n \text{ copies}},$$

where  $\text{perf}^{\text{dg}}(S, \mathcal{C}_0)$  stands for the dg enhancement of  $\text{perf}(S, \mathcal{C}_0)$  induced from  $\text{perf}_{\text{dg}}(Q)$ . Note that by Lemma 2.1, the dg category  $\text{perf}^{\text{dg}}(S, \mathcal{C}_0)$  is smooth and proper. Since the inclusion of categories  $\text{perf}(S, \mathcal{C}_0) \hookrightarrow \text{perf}(Q)$  is of Fourier–Mukai type ([23, Proposition 4.9]), its kernel  $\mathcal{K} \in \text{perf}(S \times Q, \mathcal{C}_0^{\text{op}} \boxtimes \mathcal{O}_X)$  gives rise to a Fourier–Mukai Morita equivalence

$$\Phi_{\text{dg}}^{\mathcal{K}} : \text{perf}_{\text{dg}}(S, \mathcal{C}_0) \rightarrow \text{perf}^{\text{dg}}(S, \mathcal{C}_0).$$

Hence, we can replace in the above decomposition (6.1) the dg category  $\text{perf}^{\text{dg}}(S, \mathcal{C}_0)$  by the canonical one  $\text{perf}_{\text{dg}}(S, \mathcal{C}_0)$  (see Section 2.2). Finally, using the above description (5.2) of the noncommutative nilpotence conjecture, one concludes that

$$(6.2) \quad V_{\text{NC}}(\text{perf}_{\text{dg}}(Q)) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(S, \mathcal{C}_0)) + V_{\text{NC}}(\text{perf}_{\text{dg}}(S)).$$

The proof follows now automatically from Theorem 1.1.

**Item (ii).** As proved by Kuznetsov in [23, Proposition 3.13],  $\text{perf}(S, \mathcal{C}_0)$  is Fourier–Mukai equivalent to  $\text{perf}(\tilde{S}, \mathcal{B}_0)$ . Hence, the above equivalence (6.2) reduces to

$$(6.3) \quad V_{\text{NC}}(\text{perf}_{\text{dg}}(Q)) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\tilde{S}, \mathcal{B}_0)) + V_{\text{NC}}(\text{perf}_{\text{dg}}(S)).$$

Since  $\mathcal{B}_0$  is a sheaf of Azumaya algebras and  $F$  is of characteristic zero, the canonical dg functor  $\text{perf}_{\text{dg}}(\tilde{S}) \rightarrow \text{perf}_{\text{dg}}(\tilde{S}, \mathcal{B}_0)$  becomes an isomorphism in  $\text{NChow}(k)_F$ ; see [34, Theorem 2.1]. Consequently, Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\tilde{S}, \mathcal{B}_0))$  reduces to Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(\tilde{S}))$ . The proof follows now from Theorem 1.1.

**Item (iii).** As proved by Kuznetsov in [23, Proposition 3.15],  $\text{perf}(S, \mathcal{C}_0)$  is Fourier–Mukai equivalent to  $\text{perf}(\hat{S}, \mathcal{B}_0)$ . Hence, the above equivalence (6.2) reduces to

$$(6.4) \quad V_{\text{NC}}(\text{perf}_{\text{dg}}(Q)) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\hat{S}, \mathcal{B}_0)) + V_{\text{NC}}(\text{perf}_{\text{dg}}(S)).$$

The proof of the first claim follows now from Theorem 1.1.

Let us now prove the second claim, which via (6.4) is equivalent to the proof of Conjecture  $V_{\text{NC}}(\text{perf}_{\text{dg}}(Q))$ . Thanks to Vial [35, Theorem 4.2 and Corollary 4.4], the rational Chow motive  $M_{\mathbb{Q}}(Q)$  of  $Q$  decomposes as  $M_{\mathbb{Q}}(Q) = M_{\mathbb{Q}}(S)^{\oplus(n-\dim(S))} \oplus N$ , where  $N$  stands for a submotive of a smooth projective  $k$ -scheme of dimension  $\leq \dim(S)$ . Therefore, when  $\dim(S) \leq 2$ , Conjecture  $V(Q) = V_{\text{NC}}(\text{perf}_{\text{dg}}(Q))$  holds.

**Remark 6.5.** Assume that  $S$  is a smooth projective curve and that  $k$  is algebraically closed. In this remark we provide a “categorical” proof of the second claim of item (iii) of Theorem 1.2. Thanks to the work of Graber–Harris–Starr [13], the fibration  $q : Q \rightarrow S$  admits

a section. Making use of it, we can perform reduction by hyperbolic splitting in order to obtain a conic bundle  $q' : Q' \rightarrow S$ ; consult [5, Section 1.3] for details. The sheaf  $\mathcal{C}'_0$  of even parts of the associated Clifford algebra is such that the categories  $\text{perf}(S, \mathcal{C}_0)$  and  $\text{perf}(S, \mathcal{C}'_0)$  are Fourier–Mukai equivalent; see [5, Remark 1.8.9]. As a consequence, using (6.2) and the fact that  $\dim(S) = 1$ , we obtain the following equivalence:

$$(6.6) \quad V_{\text{NC}}(\text{perf}_{\text{dg}}(Q)) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(Q')).$$

Since  $S$  is a curve,  $Q'$  is a surface. Therefore, conjecture (6.6) holds.

### 7. Proof of Theorem 1.4

**Item (i).** As proved by Kuznetsov in [23, Theorem 5.5], we have a Fourier–Mukai equivalence  $\text{perf}(X) \simeq \text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0)$  when  $m - 2r + 1 = 0$ , the semi-orthogonal decomposition  $\text{perf}(X) = \langle \text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0), \mathcal{O}_X(1), \dots, \mathcal{O}_X(m - 2r + 1) \rangle$  when  $m - 2r + 1 > 0$ , and a dual semi-orthogonal decomposition of  $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0)$  (containing a copy of  $\text{perf}(X)$  and exceptional objects) when  $m - 2r + 1 < 0$ . The proof of the case  $m - 2r + 1 = 0$  is clear. Let us now prove the case  $m - 2r + 1 > 0$ ; the proof of the case  $m - 2r + 1 < 0$  is similar. Using [8, Proposition 3.11], one obtains the following direct sum decomposition in  $\text{NChow}(k)_F$ :

$$\text{perf}_{\text{dg}}(X) \simeq \text{perf}^{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0) \oplus \underbrace{\text{perf}_{\text{dg}}(\underline{k}) \oplus \dots \oplus \text{perf}_{\text{dg}}(\underline{k})}_{(m-2r+1) \text{ copies}}.$$

Thanks to Lemma 2.1, the dg category  $\text{perf}^{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0)$  is smooth and proper.<sup>2)</sup> Since the inclusion  $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0) \hookrightarrow \text{perf}(X)$  is of Fourier–Mukai type (see [23, Proposition 4.9]), an argument similar to the one of the proof of Theorem 1.2 (i) shows us that

$$V_{\text{NC}}(\text{perf}_{\text{dg}}(X)) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(\mathbb{P}^{r-1}, \mathcal{C}_0)) + V_{\text{NC}}(\text{perf}_{\text{dg}}(\underline{k})).$$

The proof follows now automatically from Theorem 1.1.

**Items (ii)–(iii).** The proofs are similar to those of items (ii)–(iii) of Theorem 1.2.

### 8. Proof of Theorems 1.7 and 1.10

Assume that  $X_L$  and  $Y_L$  are as in classes of (i)–(ii) of Theorem 1.7 (resp. as in class (i) of Theorem 1.10). As proved by Kuznetsov in [21, Sections 10–11] (resp. in [7, Theorem 3.4]), one of the following three situations occurs:

- (a) there is a semi-orthogonal decomposition  $\text{perf}(X_L) = \langle \text{perf}(Y_L), \mathcal{E}_1, \dots, \mathcal{E}_n \rangle$ , where  $\mathcal{E}_i$  are exceptional bundles on  $X_L$ ,
- (b) there is a semi-orthogonal decomposition  $\text{perf}(Y_L) = \langle \text{perf}(X_L), \mathcal{E}'_1, \dots, \mathcal{E}'_n \rangle$ , where  $\mathcal{E}'_i$  are exceptional bundles on  $Y_L$ ,
- (c) there is a Fourier–Mukai equivalence between  $\text{perf}(X_L)$  and  $\text{perf}(Y_L)$ .

<sup>2)</sup> In the case  $m - 2r + 1 < 0$ , these properties follow from the existence of a fully faithful Fourier–Mukai functor  $\text{perf}(\mathbb{P}^{r-1}, \mathcal{C}_0) \rightarrow \text{perf}(Q)$ , with  $Q \subset \mathbb{P}^{r-1} \times \mathbb{P}^m$  a smooth hypersurface.

Therefore, the equivalence of  $V(X_L)$  and  $V(Y_L)$  is clear in situation (c). Since the inclusions of categories  $\text{perf}(Y_L) \hookrightarrow \text{perf}(X_L)$  (situation (a)) and  $\text{perf}(X_L) \hookrightarrow \text{perf}(Y_L)$  (situation (b)) are of Fourier–Mukai type, a proof similar to the one of Theorem 1.2 (i) shows us that the equivalence of  $V(X_L)$  and  $V(Y_L)$  also holds in situations (a)–(b). Note that this concludes the proof of Theorem 1.10 since Conjecture  $V(X_L)$  (resp.  $V(Y_L)$ ) holds when  $\dim(X_L) \leq 2$  (resp.  $\dim(Y_L) \leq 2$ ).

Let us now focus on class (i) of Theorem 1.7. The smooth projective  $k$ -schemes  $X_L$  and  $Y_L$  are of dimensions  $8 - r$  and  $r - 2$ , respectively. Hence,  $V(Y_L)$  holds when  $r \leq 4$  and  $V(X_L)$  when  $r = 6$ . When  $r = 5$ ,  $X_L$  (and  $Y_L$ ) is a Fano 3-fold. As explained by Gorchinskiy and Guletskii in [12, Section 5], the Chow motive of  $X$  admits a decomposition into Lefschetz motives and submotives of curves. This implies that  $V(X_L)$  also holds.

Let us now focus on class (ii) of Theorem 1.7. The smooth projective  $k$ -schemes  $X_L$  and  $Y_L$  are of dimensions  $10 - r$  and  $r - 4$ , respectively. Hence,  $V(Y_L)$  holds when  $r \leq 6$  and  $V(X_L)$  when  $r \geq 8$ . This achieves the proof.

### 9. Proof of Theorem 1.12

Following Kuznetsov [22, Section 4], let us denote by  $\alpha_i$  the orthogonal complement of  $\mathbb{A}_{i+1}$  in  $\mathbb{A}_i$ ; these are called the “primitive subcategories” in [22, Section 4]. Since Conjecture  $V_{\text{NC}}(\mathbb{A}_0^{\text{dg}})$  holds, we hence conclude by induction that Conjectures  $V_{\text{NC}}(\mathbb{A}_i^{\text{dg}})$  and  $V_{\text{NC}}(\alpha_i^{\text{dg}})$  also hold for every  $i$ . Thanks to HPD (see [22, Theorem 6.3]),  $\text{perf}(Y)$  admits a Lefschetz decomposition  $\text{perf}(Y) = \langle \mathbb{B}_m(-m), \dots, \mathbb{B}_0 \rangle$  with respect to  $\mathcal{O}_Y(1)$  such that the primitive subcategories  $\mathfrak{b}_i$  coincide (via a Fourier–Mukai functor) with the primitive subcategories  $\alpha_i$ . Consequently,  $V_{\text{NC}}(\mathfrak{b}_i^{\text{dg}})$  holds for every  $i$ . An inductive argument, starting with  $\mathfrak{b}_m = \mathbb{B}_m$ , allows us then to conclude that Conjecture  $V_{\text{NC}}(\mathbb{B}_i^{\text{dg}})$  also holds for any  $i$ . Now, thanks once also to HPD (see [22, Thm. 5.3]), there exists also a triangulated category  $\mathbb{C}_L$  and semi-orthogonal decompositions

$$\begin{aligned} \text{perf}(X_L) &= \langle \mathbb{C}_L, \mathbb{A}_{\dim(L)}(1), \dots, \mathbb{A}_n(n - \dim(L)) \rangle, \\ \text{perf}(Y_L) &= \langle \mathbb{B}_m(\dim(L^\perp) - m), \dots, \mathbb{B}_{\dim(L^\perp)}(-1), \mathbb{C}_L \rangle. \end{aligned}$$

Moreover, the composed functor  $\text{perf}(X_L) \rightarrow \mathbb{C}_L \rightarrow \text{perf}(Y_L)$  is of Fourier–Mukai type. As a conclusion, since  $X_L$  and  $Y_L$  are smooth, we can apply Theorem 1.1 and obtain the following chain of equivalences:

$$V(X_L) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(X_L)) \iff V_{\text{NC}}(\mathbb{C}_L^{\text{dg}}) \iff V_{\text{NC}}(\text{perf}_{\text{dg}}(Y_L)) \iff V(Y_L).$$

This achieves the proof.

### 10. Proof of Theorem 1.14

Thanks to the work of Cossec [11], the conic bundle  $Y \rightarrow \mathbb{P}^2$  has a natural structure of quartic double solid  $Y \rightarrow \mathbb{P}^3$  ramified along a quartic symmetroid  $D$ . Via the natural involution on the resolution of singularities of  $D$ , one hence obtains an Enriques surface  $S$ ; consult [11, Section 3] for details. As proved by Zube in [38, Section 5], one has moreover a semi-orthogonal decomposition  $\text{perf}(S) = \langle \mathcal{T}_S, \mathcal{E}_1, \dots, \mathcal{E}_{10} \rangle$ , where  $\mathcal{E}_i$  are exceptional objects. Let

us denote by  $\mathcal{T}_S^{\text{dg}}$  the dg enhancement of  $\mathcal{T}_S$  induced from  $\text{perf}_{\text{dg}}(S)$ . Thanks to Lemma 2.1,  $\mathcal{T}_S^{\text{dg}}$  is smooth and proper. Hence, since  $S$  is a surface, an argument similar to the one of the proof of Theorem 1.2 (i) shows us that Conjecture  $V_{\text{NC}}(\mathcal{T}_S^{\text{dg}})$  holds.

Now, recall from Ingalls and Kuznetsov [15, Section 5.5] the construction of the Fourier–Mukai functor  $\Phi : \text{perf}(S) \rightarrow \text{perf}(X)$  whose restriction to  $\mathcal{T}_S$  is fully-faithful. As proved in [15, Proposition 3.8 and Theorem 4.3], one has a semi-orthogonal decomposition

$$\text{perf}(X) = \langle \Phi(\mathcal{T}_S), \mathcal{E}'_1, \mathcal{E}'_2 \rangle,$$

where  $\mathcal{E}'_i$  are exceptional objects. As a consequence, we obtain the equivalence

$$(10.1) \quad V_{\text{NC}}(\text{perf}_{\text{dg}}(X)) \iff V_{\text{NC}}(\Phi(\mathcal{T}_S)^{\text{dg}}),$$

where  $\Phi(\mathcal{T}_S)^{\text{dg}}$  stands for the dg enhancement of  $\Phi(\mathcal{T}_S)$  induced from  $\text{perf}_{\text{dg}}(X)$ . Since the kernel  $\mathcal{K}$  of the above Fourier–Mukai functor  $\Phi$  gives rise to a Morita equivalence

$$\Phi_{\text{dg}}^{\mathcal{K}} : \mathcal{T}_S^{\text{dg}} \rightarrow \Phi(\mathcal{T}_S)^{\text{dg}},$$

we conclude that conjecture (10.1) also holds. This achieves the proof.

**Acknowledgement.** The authors are grateful to Bruno Kahn and Claire Voisin for useful comments and answers, as well as to the anonymous referee for useful remarks.

## References

- [1] *Y. André*, Une introduction aux motifs (motifs purs, motifs mixtes, périodes), Panor. Synthèses **17**, Société Mathématique de France, Paris 2004.
- [2] *Y. André* and *B. Kahn*, Nilpotence, radicaux et structures monoïdales, Rend. Semin. Mat. Univ. Padova **108** (2002), 107–291.
- [3] *M. Artin*, Algebraization of formal moduli. II: Existence of modification, Ann. of Math. (2) **91** (1970), 88–135.
- [4] *M. Artin* and *D. Mumford*, Some elementary examples of unirational varieties which are not rational, Proc. Lond. Math. Soc. (3) **25** (1972), 75–95.
- [5] *A. Auel*, *M. Bernardara* and *M. Bolognesi*, Fibrations in complete intersections of quadrics, Clifford algebras, derived categories, and rationality problems, J. Math. Pures Appl. (9) **102** (2014), 249–291.
- [6] *H. Bass*, Algebraic K-theory, W. A. Benjamin, New York 1968.
- [7] *M. Bernardara*, *M. Bolognesi* and *D. Faenzi*, Homological projective duality for determinantal varieties, preprint 2014, <http://arxiv.org/abs/1410.7803>.
- [8] *M. Bernardara* and *G. Tabuada*, Chow groups of intersections of quadrics via homological projective duality and (Jacobians of) noncommutative motives, preprint 2013, <http://arxiv.org/abs/1310.6020>.
- [9] *A. Bondal* and *D. Orlov*, Derived categories of coherent sheaves, in: International congress of mathematicians. Vol. II: Invited lectures (Beijing 2002), Higher Education Press, Beijing (2002), 47–56.
- [10] *D.-C. Cisinski* and *G. Tabuada*, Symmetric monoidal structure on noncommutative motives, J. K-Theory **9** (2012), no. 2, 201–268.
- [11] *F. Cossec*, Reye congruences, Trans. Amer. Math. Soc. **280** (1983), no. 2, 737–751.
- [12] *S. Gorchinskiy* and *V. Guletskii*, Motives and representability of algebraic cycles on threefolds over a field, J. Algebraic Geom. **21** (2012), no. 2, 347–373.
- [13] *T. Graber*, *J. Harris* and *J. Starr*, Families of rationally connected varieties, J. Amer. Math. Soc. **16** (2003), no. 1, 57–67.
- [14] *D. Huybrechts*, Fourier–Mukai transforms in algebraic geometry, Oxford Math. Monogr., Clarendon Press, Oxford, 2006.
- [15] *C. Ingalls* and *A. Kuznetsov*, On nodal Enriques surfaces and quartic double solids, preprint 2010, <http://arxiv.org/abs/1012.3530>.
- [16] *B. Kahn* and *R. Sebastian*, Smash-nilpotent cycles on abelian 3-folds, Math. Res. Lett. **16** (2009), no. 6, 1007–1010.

- [17] *B. Keller*, On differential graded categories, in: International congress of mathematicians. Vol. II: Invited lectures (Madrid 2006), European Mathematical Society, Zürich (2006), 151–190.
- [18] *M. Kontsevich*, Noncommutative motives, Talk at the Institute for Advanced Study on the occasion of the 61st birthday of Pierre Deligne, Princeton, October 2005. Video available at <http://video.ias.edu/Geometry-and-Arithmetic>.
- [19] *M. Kontsevich*, Notes on motives in finite characteristic, in: Algebra, arithmetic, and geometry. In honor of Yu. I. Manin. Vol. II, Progr. Math. **270**, Birkhäuser, Boston (2009), 213–247.
- [20] *M. Kontsevich*, Mixed noncommutative motives, Talk at the workshop on homological mirror symmetry, Miami 2010. Notes available at <https://math.berkeley.edu/~auroux/frg/miami10-notes/>.
- [21] *A. Kuznetsov*, Homological projective duality for Grassmannians of lines, preprint 2006, <http://arxiv.org/abs/math/0610957>.
- [22] *A. Kuznetsov*, Homological projective duality, Publ. Math. Inst. Hautes Études Sci. **105** (2007), 157–220.
- [23] *A. Kuznetsov*, Derived categories of quadric fibrations and intersections of quadrics, Adv. Math. **218** (2008), no. 5, 1340–1369.
- [24] *A. Kuznetsov*, Semiorthogonal decompositions in algebraic geometry, preprint 2014, <http://arxiv.org/abs/1404.3143>.
- [25] *V. Lunts* and *D. Orlov*, Uniqueness of enhancement for triangulated categories, J. Amer. Math. Soc. **23** (2010), 853–908.
- [26] *M. Marcolli* and *G. Tabuada*, Unconditional motivic Galois groups and Voevodsky's nilpotence conjecture in the noncommutative world, preprint 2011, <http://arxiv.org/abs/1112.5422v1>.
- [27] *M. Marcolli* and *G. Tabuada*, Kontsevich's noncommutative numerical motives, Compos. Math. **148** (2012), no. 6, 1811–1820.
- [28] *M. Marcolli* and *G. Tabuada*, Noncommutative motives and their applications, preprint 2013, <http://arxiv.org/abs/1311.2867>; to appear in MSRI publications, Berkeley.
- [29] *M. Marcolli* and *G. Tabuada*, Noncommutative motives, numerical equivalence, and semi-simplicity, Amer. J. Math. **136** (2014), no. 1, 59–75.
- [30] *T. Matsusaka*, The criteria for algebraic equivalence and the torsion group, Amer. J. Math. **79** (1957), 53–66.
- [31] *B. G. Moishezon*, On  $n$ -dimensional compact varieties with  $n$  algebraically independent meromorphic functions. I, II and III, Amer. Math. Soc. Transl. II **63** (1967), 51–177.
- [32] *G. Tabuada*, A guided tour through the garden of noncommutative motives, Clay Math. Proc. **16** (2012), 259–276.
- [33] *G. Tabuada*, Chow motives versus noncommutative motives, J. Noncommut. Geom. **7** (2013), no. 3, 767–786.
- [34] *G. Tabuada* and *M. Van den Bergh*, Noncommutative motives of Azumaya algebras, J. Inst. Math. Jussieu **14** (2015), no. 2, 379–403.
- [35] *C. Vial*, Algebraic cycles and fibrations, Doc. Math. **18** (2013), 1521–1553.
- [36] *V. Voevodsky*, A nilpotence theorem for cycles algebraically equivalent to zero, Int. Math. Res. Not. IMRN **1995** (1995), no. 4, 187–198.
- [37] *C. Voisin*, Remarks on zero-cycles of self-products of varieties, in: Moduli of vector bundles (Sanda 1994, Kyoto 1994), Lecture Notes in Pure and Appl. Math. **179**, Dekker, New York (1996), 265–285.
- [38] *S. Zube*, Exceptional vector bundles on Enriques surfaces, Math. Notes **61** (1997), no. 6, 693–699.

---

Marcello Bernardara, Institut de Mathématiques de Toulouse, Université Paul Sabatier,  
118 route de Narbonne, 31062 Toulouse Cedex 9, France  
e-mail: marcello.bernardara@math.univ-toulouse.fr

Matilde Marcolli, Mathematics Department, Mail Code 253-37, Caltech,  
1200 E. California Blvd., Pasadena, CA 91125, USA  
e-mail: matilde@caltech.edu

Gonçalo Tabuada, Department of Mathematics, MIT, Cambridge, MA 02139, USA;  
and Departamento de Matemática, FCT, UNL, Portugal;  
and Centro de Matemática e Aplicações (CMA), FCT, UNL, Portugal  
e-mail: tabuada@math.mit.edu

Eingegangen 21. November 2014